

Product Datasheet / Instruction for use



Rabbit IgG anti-Bovine IgM (μ)-HRPO, MinX none

General Information

Catalog Number: SE	C-182490	Physical State:	Lyophilized	
Quantity:	20 mg	Reconstitution Volume:	2.0 mL	
		Reconstitution Buffer:	Restore with deionized water (or equivalent) at room temperature	
		Shipping Conditions:		
		Product Expiration	Expiration date is one (1) year from date of opening.	
Antibody Host / Forn	nat	I		
Host Species: Ra	bbit	Clonality / Isotype / Clon	e: polyclonal Ig	
Antibody Format: Ig0	G whole molecule	Concentration:	10.0 mg/mL by UV absorbance at 280 nm	
		Conjugation:	Horseradish Peroxidase (HRP)	
		Maximum Excitation / Emission: /		
Specificity		I		
Target Species:	Bovine	Immunogen:	Bovine IgM mu heavy chain	
Antibody Specificity:	lgM (μ-Chain/ Fc Fragment)	This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by cross adsorption and extensive dialysis against the buffer stated above. Assa by immunoelectrophoresis resulted in a single precipitin arc against ant Peroxidase, anti-Rabbit Serum, Bovine IgM and Bovine Serum. No reaction was observed against Bovine IgG.		
		Minimal Cross Reactivity: none		
Formulation, Transpo	ort and Storage	1		
· · · · · · · · · · · · · · · · · · ·	Storage Buffer:0.02 M PotassiumPhosphate, 0.15 M Sodium Chloride, pH 7.2Stabilizer:10 mg/ml Polyethylene			
Phosphate, 0.15 M Sodi Stabilizer: 10 mg	ium Chloride, pH 7.2	contents and freeze at -2 thawing. Centrifuge proc	restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and duct if not completely clear after standing at product is stable for several weeks at 4° C as a	
Phosphate, 0.15 M Sodi Stabilizer: 10 mg Glycol (PEG-8000)	ium Chloride, pH 7.2 ;/ml Polyethylene	contents and freeze at -2 thawing. Centrifuge proc room temperature. This	0° C or below. Avoid cycles of freezing and duct if not completely clear after standing at	
Phosphate, 0.15 M Sodi Stabilizer: 10 mg Glycol (PEG-8000)	ium Chloride, pH 7.2	contents and freeze at -2 thawing. Centrifuge proc room temperature. This	0° C or below. Avoid cycles of freezing and duct if not completely clear after standing at product is stable for several weeks at 4° C as a	

This product is for In Vitro exp	erimental use only. Not for Ther	apeutic or Diagnostic use.		
Manufacturer				Management System
BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771



Product Datasheet / Instruction for use



Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

Dilution: ELISA 1:10.000 - 1:50.000, Immunohistochemistry (IHC) 1:500 - 1:2,500, Western Blot (WB) 1:1.000 - 1:10.000

Application Note:

This product has been assayed against 1.0 ug of Bovine IgM in a standard capture ELISA using ABTS (2,2'-azino-bis-[3ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000 of the reconstitution concentration is suggested for this product.

Background Information

Immunoglobulin M is the largest antibody isotype and the first to be secrected against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together, the approixmate molecular weight of IgM is 900kDa and possesses 10 binding sites (though due to the size of most antigens, not all sites are capable of binding at once). Due to this large size, IgM is typically isolated to the serum. Anti-Bovine IgM antibody is ideal for investigators in Immunology, Microbiology, and Cell Biology.

Safety Information

This reagent contains Gentamicin Sulfate (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.DO NOT ADD SODIUM AZIDE! Use of Sodium Azide as a preservative will substantially inhibit the enzyme activity of horseradish peroxidase!

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL **Diagnostica Vertrieb GmbH** Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL



Management System ISO 9001:2015