

Product Datasheet / Instruction for use



Rabbit IgG anti-Dog IgM (μ)-HRPO, MinX none

General Information

Catalog Number: SEC	C-182517	Physical State:	Lyophilized		
Quantity:	20 mg	Reconstitution Volume:	2.0 mL		
		Reconstitution Buffer:	Restore with deionized water (or equivalent)		
		Shipping Conditions:	at room temperature		
		Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / Form	nat				
Host Species: Ral	bbit	Clonality / Isotype / Clon	e: polyclonal Ig		
Antibody Format: IgG	whole molecule	Concentration:	10.0 mg/mL by UV absorbance at 280 nm		
		Conjugation:	Horseradish Peroxidase (HRP)		
		Maximum Excitation / Emission: /			
Specificity					
Target Species:	Dog	Immunogen:	Dog IgM mu heavy chain		
Antibody Specificity:	lgM (μ-Chain/ Fc Fragment)	Dog IgM mu chain Antibody Peroxidase Conjugated is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin ar against anti-Peroxidase, anti-Rabbit Serum, Dog IgM and Dog Serum. No reaction was observed against Dog IgG.			
		Minimal Cross Reactivity: none			
Formulation, Transpo	ort and Storage				
Storage Buffer:0.02 M PotassiumPhosphate, 0.15 M Sodium Chloride, pH 7.2Stabilizer:10 mg/mL Bovine SerumAlbumin (BSA) - Immunoglobulin and Protease		Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.			
free Preservatives: 0.01% (w/v) Gentamicin Sulfate		Storage Temperature: 4-8°C			
Sulfate	、 <i>, ,</i>				

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.						
Manufacturer				Management System		
BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015		
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771		



Product Datasheet / Instruction for use



Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

Dilution: ELISA 1:10.000 - 1:50.000, Immunohistochemistry (IHC) 1:500 - 1:2,500, Western Blot (WB) 1:1.000 - 1:10.000

Application Note:

Dog IgM mu chain Antibody Peroxidase Conjugated has been assayed against 1.0 ug of Dog IgM in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.

Background Information

Anti-Dog IgM antibody specifically detects dog IgM. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together, the approximate molecular weight of IgM is 900kDa and possesses 10 binding sites (though due to the size of most antigens, not all sites are capable of binding at once). Due to this large size, IgM is typically isolated to the serum. Anti-Dog IgM antibody is ideal for investigators in Immunology, Microbiology, and Cell Biology.

Safety Information

This reagent contains Gentamicin Sulfate (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.DO NOT ADD SODIUM AZIDE! Use of Sodium Azide as a preservative will substantially inhibit the enzyme activity of horseradish peroxidase!

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL **Diagnostica Vertrieb GmbH** Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL



Management System ISO 9001:2015