

Product Datasheet / Instruction for use



# Rabbit IgG anti-Human IgG (Fc)-FITC, MinX none

# **General Information**

Catalog Number: SEC-182578	Physical State:	Lyophilized		
Quantity: 20 mg	Reconstitution Volume:	2.0 mL		
	<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)		
	Shipping Conditions:	at room temperature		
	Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / Format	1			
Host Species: Rabbit	Clonality / Isotype / Clon	ne: polyclonal Ig		
Antibody Format: IgG whole molecule	Concentration:	10.0 mg/mL by UV absorbance at 280 nm		
	Conjugation:	FITC (Fluoresceinisothiocyanat)		
	Maximum Excitation / Emission: 492 nm / 520 nm			
Specificity				
Target Species: Human	Immunogen:	Human IgG F(c) fragment		
Antibody Specificity: IgG (Fc Fragment)	This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophores resulted in a single precipitin arc against anti-fluorescein, anti-Rabbit Serum, Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against Human IgG F(ab).			
	Minimal Cross Reactivity: none			
	Minimal Cross Reactivity	: none		
Formulation, Transport and Storage	Minimal Cross Reactivity	: none		
Storage Buffer: 0.01 M Sodium Phosphate,	Store vial at 4° C prior to contents and freeze at -2	restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and		
Formulation, Transport and StorageStorage Buffer:0.01 M Sodium Phosphate,0.15 M Sodium Chloride, pH 7.2Stabilizer:10 mg/mL Bovine SerumAlbumin (BSA) - Immunoglobulin and Proteasefree	Store vial at 4° C prior to contents and freeze at -2 thawing. Centrifuge proc room temperature. This undiluted liquid. Dilute o	restoration. For extended storage aliquot		
Storage Buffer:0.01 M Sodium Phosphate,0.15 M Sodium Chloride, pH 7.2Stabilizer:10 mg/mL Bovine SerumAlbumin (BSA) - Immunoglobulin and Protease	Store vial at 4° C prior to contents and freeze at -2 thawing. Centrifuge proc room temperature. This	restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and luct if not completely clear after standing at product is stable for several weeks at 4° C as ar		

This product is for In Vitro exp	erimental use only. Not for Ther	apeutic or Diagnostic use.			
Manufacturer				Management System	
<b>BIOZOL</b> Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015	
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771	



#### Product Datasheet / Instruction for use



Application: FLISA, Flow Cytometry, Immunofluorescence

FLISA 1:10.000 - 1:50.000, Flow cytometry 1:500 - 1:2,500, Immunofluorescence 1:1,000 - 1:5,000 Dilution:

#### **Application Note:**

Anti-Human IgG F(c) Fluorescein Conjugated is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

### **Background Information**

Anti-Human IgG F(c) Fluorescein Conjugated generated in rabbit detects Human F(c). A proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of human IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(c) Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

# Safety Information

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

# Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL **Diagnostica Vertrieb GmbH** Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL



Management System ISO 9001:2015