

## Goat IgG anti-Guinea Pig IgG (H+L)-unconj., MinX Bo,Ck,Go,Hm,Ho,Hu,Ms,Rb,Rt,Sh

### General Information

<b>Catalog Number:</b> SEC-182897	<b>Physical State:</b> Liquid (sterile filtered)
<b>Quantity:</b> 1 mg	<b>Reconstitution Volume:</b>
	<b>Reconstitution Buffer:</b>
	<b>Shipping Conditions:</b> Wet Ice
	<b>Product Expiration:</b> Expiration date is one (1) year from date of opening.

### Antibody Host / Format

<b>Host Species:</b> Goat	<b>Clonality / Isotype / Clone:</b> polyclonal Ig
<b>Antibody Format:</b> IgG whole molecule	<b>Concentration:</b> 1.0 mg/mL by UV absorbance at 280 nm
	<b>Conjugation:</b> unconjugated
	<b>Maximum Excitation / Emission:</b> /


### Specificity

<b>Target Species:</b> Guinea Pig	<b>Immunogen:</b> Guinea Pig IgG whole molecule
<b>Antibody Specificity:</b> IgG (H+L)	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Guinea Pig IgG and Guinea Pig Serum. No reaction was observed against Bovine, Chicken, Goat, Hamster, Horse, Human, Mouse, Rabbit, Rat and Sheep Serum Proteins. Minimal cross reactivity was observed against the species listed above.
	<b>Minimal Cross Reactivity:</b> Bovine, Chicken, Goat, Hamster generally, Horse, Human, Mouse, Rabbit, Rat, Sheep

### Formulation, Transport and Storage

<b>Storage Buffer:</b> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
<b>Stabilizer:</b> None	
<b>Preservatives:</b> 0.01% (w/v) Sodium Azide	
	<b>Storage Temperature:</b> 4-8°C

### Application Recommendation

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.		
<b>Manufacturer</b> <b>BIOZOL</b> Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de	<a href="http://www.dianova.com">www.dianova.com</a> Your Secondary Antibody Portal Powered by BIOZOL
	 TÜVRheinland ZERTIFIZIERT	Management System ISO 9001:2015 www.tuv.com ID 900019771

**Application:** ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

**Dilution:** ELISA 1:50.000, Immunohistochemistry (IHC) 1:1,000 - 1:5,000, Western Blot (WB) 1:2.000 - 1:10.000

**Application Note:**

Anti-Guinea Pig IgG antibody is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency.

**Background Information**

Anti-Guinea Pig IgG Antibody generated in goat detects guinea pig IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsinization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

**Safety Information**

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

**Disclaimer**

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

**Manufacturer**

**BIOZOL**  
Diagnostica Vertrieb GmbH  
Leipziger Straße 4  
85386 Eching

Phone +49 (89) 3799 666 6  
Fax +49 (89) 3799 666 99  
E-mail info@biozol.de  
www.biozol.de

[www.dianova.com](http://www.dianova.com)

Your Secondary Antibody Portal  
Powered by BIOZOL



Management System  
ISO 9001:2015



www.tuv.com  
ID 900019771