

Product Datasheet / Instruction for use



Goat IgG anti-Mouse IgG2a (Fc)-ATTO 425, MinX none

General Information

Catalog Number: SE	C-183183	Physical State: Lyophilized		
Quantity:	0,5 mg	Reconstitution Volume:	500 μL	
		Reconstitution Buffer:	Restore with deionized water (or equivalent) at room temperature	
		Shipping Conditions:		
		Product Expiration	Expiration date is one (1) year from date of opening.	
Antibody Host / Forn	mat			
Host Species: Go	pat	Clonality / Isotype / Clone: polyclonal Ig		
Antibody Format: Ig	G whole molecule	Concentration:	1.0 mg/mL by UV absorbance at 280 nm	
		Conjugation:	ATTO 425	
		Maximum Excitation / Emission: 439 nm / 485 nm		
Specificity				
Target Species:	Mouse	Immunogen:	Mouse IgG2a heavy chain	
Antibody Specificity:	lgG2a (lgG Fc gamma 2a)	vas prepared from monospecific antiserum by ography using antigens coupled to agarose bead		
		reactivities. Assay by imp precipitin arc against anti Specificity was confirmed other mouse heavy or ligh against Bovine, Human, a	dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a.	
		reactivities. Assay by imp precipitin arc against anti Specificity was confirmed other mouse heavy or ligh against Bovine, Human, a	dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a. by ELISA at less than 1% cross-reactivity agains at chain isotypes. No reaction was observed nd Rabbit Serum Proteins. Specificity was s than 1% of target signal.	
Formulation, Transpo		reactivities. Assay by imp precipitin arc against anti Specificity was confirmed other mouse heavy or ligh against Bovine, Human, a confirmed by ELISA at less	dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a. by ELISA at less than 1% cross-reactivity agains at chain isotypes. No reaction was observed nd Rabbit Serum Proteins. Specificity was s than 1% of target signal.	
Storage Buffer: 0.02 M Phosphate, 0.15 M Sod	ort and Storage VI Potassium ium Chloride, pH 7.2	reactivities. Assay by imprecipitin arc against anti Specificity was confirmed other mouse heavy or ligh against Bovine, Human, a confirmed by ELISA at less Minimal Cross Reactivity Store vial at 4° C prior to a contents and freeze at -20	dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a. by ELISA at less than 1% cross-reactivity agains nt chain isotypes. No reaction was observed nd Rabbit Serum Proteins. Specificity was s than 1% of target signal.	
Storage Buffer: 0.02 M Phosphate, 0.15 M Sod Stabilizer: 10 mg	ort and Storage VI Potassium	reactivities. Assay by imprecipitin arc against anti Specificity was confirmed other mouse heavy or ligh against Bovine, Human, a confirmed by ELISA at less Minimal Cross Reactivity Store vial at 4° C prior to a contents and freeze at -20 thawing. Centrifuge prod room temperature. This undiluted liquid. Dilute o	dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a. by ELISA at less than 1% cross-reactivity agains at chain isotypes. No reaction was observed nd Rabbit Serum Proteins. Specificity was s than 1% of target signal. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and luct if not completely clear after standing at product is stable for several weeks at 4° C as ar nly prior to immediate use.	
Storage Buffer: 0.02 M Phosphate, 0.15 M Sod Stabilizer: 10 mg Albumin (BSA) - Immun free	ort and Storage M Potassium lium Chloride, pH 7.2 g/mL Bovine Serum	reactivities. Assay by imprecipitin arc against anti Specificity was confirmed other mouse heavy or ligh against Bovine, Human, a confirmed by ELISA at less Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge prod room temperature. This	dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a. by ELISA at less than 1% cross-reactivity agains nt chain isotypes. No reaction was observed nd Rabbit Serum Proteins. Specificity was s than 1% of target signal. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and luct if not completely clear after standing at product is stable for several weeks at 4° C as an	

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.						
Manufacturer				Management System		
BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015		
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771		



Product Datasheet / Instruction for use



Application: FLISA, Flow Cytometry, Immunofluorescence, Western Blot

Dilution: FLISA >1:20.000, Flow cytometry 1:500 - 1:2,500, Immunofluorescence >1:5,000, Western Blot (WB) 1:4.000 - 1:20.000

Application Note:

Mouse IgG2a secondary antibody is available in a variety of formats. ATTO 425 conjugations are designed for STED microscopy, FRET, immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Anti-Mouse IgG2a secondary antibody is suitable for ELISA, Immunohistochemistry western blotting as well as other anti IgG2a antibody based assays. The emission spectra for this ATTO conjugate matches the principle output wavelengths of most common fluorescence instrumentation.

Background Information

Anti-Mouse IgG2a ATTO 425 Antibody generated in goat detects reactivity to Mouse IgG2a (Gamma 2a chain). Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. IgG2, the second largest of IgG isotypes, comprises almost 25% of IgG and has a low affinity for binding to the Fc receptor of phagocytic cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Safety Information

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT Www.tuv.com ID 900019771

Management System ISO 9001:2015