

Donkey F(ab')₂ anti-Human IgM (μ)-RPE, MinX Bo, Ho

General Information

| | |
|-----------------------------------|--|
| Catalog Number: SEC-183767 | Physical State: Lyophilized |
| Quantity: 1 ml | Reconstitution Volume: 1.0 mL |
| | Reconstitution Buffer: Restore with deionized water (or equivalent) |
| | Shipping Conditions: at room temperature |
| | Product Expiration: Expiration date is six (6) months from date of opening. |

Antibody Host / Format

| | |
|--|---|
| Host Species: Donkey | Clonality / Isotype / Clone: polyclonal Ig |
| Antibody Format: IgG F(ab') ₂ Fragment | Concentration: 0.5 mg/mL by absorbance = 82.0 at 565 nm |
| | Conjugation: RPE (R-phycoerythrin) |
| | Maximum Excitation / Emission: 490 nm; 545 nm; 566 nm / 580 nm |

Specificity

| | |
|---|---|
| Target Species: Human | Immunogen: Human IgM Fc ₅ μ fragment |
| Antibody Specificity: IgM (μ-Chain/ Fc Fragment) | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Phycoerythrin, anti-Donkey Serum, Human IgM, Human IgM Fc ₅ μ and Human Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c), Bovine or Horse Serum Proteins, or other human heavy or light chain proteins. |
| | Minimal Cross Reactivity: Bovine, Horse |

Formulation, Transport and Storage

| | |
|---|---|
| Storage Buffer: 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2 | Store vial at 4° C prior to restoration. Restore with deionized water (or equivalent). This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Centrifuge product if not completely clear after standing at room temperature. Do not freeze after reconstitution. Store reagent in the dark. Use subdued lighting during handling and incubation of cells prior to analysis. |
| Stabilizer: 15 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free | |
| Preservatives: 0.05% (w/v) Sodium Azide | |
| | Storage Temperature: 4-8°C |

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL
Diagnostica Vertrieb GmbH
Leipziger Straße 4
85386 Eching

Phone +49 (89) 3799 666 6
Fax +49 (89) 3799 666 99
E-mail info@biozol.de
www.biozol.de

www.dianova.com

Your Secondary Antibody Portal
Powered by BIOZOL



Management System
ISO 9001:2015

www.tuv.com
ID 900019771



Application Recommendation

Application: Flow Cytometry, Immunofluorescence

Dilution: Flow cytometry 1:100 - 1:250, Immunofluorescence 1:100 - 1:250

Application Note:

F(ab')₂ Anti-Human IgM Fc₅μ Phycoerythrin Antibody is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10⁶ cells in flow cytometry is approximately 1.0 μg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:50 to 1:200 should be suitable for most applications.

Background Information

F(ab')₂ Anti-Human IgM Fc₅μ Phycoerythrin Antibody generated in donkey detects specifically the Fc₅μ portion of the human IgM heavy chain. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together. F(ab')₂ Anti-Human IgM Fc₅μ antibody is ideal for investigators in Immunology, Microbiology, and Cell Biology.

Safety Information

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL
Diagnostica Vertrieb GmbH
Leipziger Straße 4
85386 Eching

Phone +49 (89) 3799 666 6
Fax +49 (89) 3799 666 99
E-mail info@biozol.de
www.biozol.de

www.dianova.com

Your Secondary Antibody Portal
Powered by BIOZOL



Management System
ISO 9001:2015



www.tuv.com
ID 900019771