



Collagen II Ab-1 (Clone 5B2.5)

Mouse Monoclonal Antibody

Cat. #DLN-08487, DLN-08488, or DLN-08486 (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified Ab with BSA and Azide)

Cat. #DLN-08489 or DLN-08490 (0.1ml or 0.2ml at 1.0mg/ml) (Purified Ab without BSA and Azide)

Comments: Ab-1 is highly specific to type II collagen and shows no cross-reaction with types I, III, IV, V, VI, IX, X, or XI. In Western blotting, 5B2.5 MAb reacts with the TC_A fragment of lathyritic type II collagen after digestion with mammalian collagenase. It does not react with pepsin-digested type II collagen.

Epitope: GGFDEK, slightly preceding the lysine residue involved in crosslinking. Two terminal glycine residues constitute a key structural element of the epitope. The epitope is lost after digestion of lathyritic type II collagen with recombinant stromelysin.

Species Reactivity: Human, Chicken, and Mouse. Others-not known

Clone Designation: 5B2.5

Ig Isotype / Light Chain: IgG_{2a} / |

Immunogen: A purified preparation of lathyritic type II collagen from embryonic chicken sternum

Applications and Suggested Dilutions:

- Immunofluorescence
- Western Blotting¹ (Ab 1-2µg/ml for 2hrs at RT)
- Immunohistology (Not suitable)

The optimal dilution for a specific application should be determined by the investigator.

Positive Control: Cartilage

Cellular Localization: Cartilage matrix

Supplied As:

200µg/ml antibody purified by Protein A chromatography. Prepared in 10mM PBS, pH 7.4, with 0.2% BSA and 0.09% sodium azide. Also available without BSA and azide at 1mg/ml.

Storage and Stability: Ab with sodium azide is stable for 24 months when stored at 2-8°C. Antibody WITHOUT sodium azide is stable for 36 months when stored at below 0°C.

Key References:

1. Mayne R; et al. Connective Tissue Research, 1994, 31(1):11- 21.
2. Chen Q; et al. Developmental Dynamics, 1993 Jan, 196(1):47-53.

dianova GmbH

Warburgstr. 45 • 20354 Hamburg

Telefon (040)45067-0 • Telefax (040) 45067-490 • www.dianova.de



Collagen II Ab-1 (Clone 5B2.5)

Mouse Monoclonal Antibody

Cat. #DLN-08487, DLN-08488, or DLN-08486 (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified Ab with BSA and Azide)

Cat. #DLN-08489 or DLN-08490 (0.1ml or 0.2ml at 1.0mg/ml) (Purified Ab without BSA and Azide)

Limitations and Warranty:

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the actual price paid for the product. Dianova is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

Material Safety Data:

This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.

For Research Use Only

dianova GmbH

Warburgstr. 45 • 20354 Hamburg

Telefon (040)45067-0 • Telefax (040) 45067-490 • www.dianova.de