

DATA SHEET

CD1a Ab-5 (Clone O10)

Mouse Monoclonal Antibody

Catalog #DLN-14729, DLN-14730, or DLN-14733 (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) Cat. #DLN-14731 or DLN-14734 (0.1ml or 0.2ml at 1.0mg/ml) (Purified Ab without BSA and Azide) Catalog #DLN-14732 (7.0ml)

Description: At least five CD1 genes (CD1a, b, c, d, and e) are identified. CD1 is expressed on cortical thymocytes, Langerhans cells, and dendritic cells. It is absent on mature peripheral blood T cells but intracytoplasmic expression is detected on activated T lymphocytes. CD1 proteins have been demonstrated to restrict T-cell response to non-peptide lipid and glycolipid antigens and play a role in non-classical antigen presentation.

Comments: Ab-5 detects cortical thymocytes, Langerhans cells in epidermis, dendritic cells of dermis and Langherhans cells of mucosa of tonsil. It may also detect small focal groups of lymphocytes outside the germinal centers of tonsil indicating a crossreaction with CD1b. This antibody is useful in the characterization of leukaemias and lymphomas.

Mol. Wt. of Antigen: 49kDa

Species Reactivity: Human. Others not-tested.

Clone Designation: 010

Ig Isotype / Light Chain: IgG₁/κ

Applications and Suggested Dilutions:

- Immunohistology (Formalin/paraffin) (Use Ab at 1:50-1:100 for 30 min at RT)
- * (Staining of formalin-fixed tissues REQUIRES boiling tissue sections in 1mM EDTA, pH 8.0, for 10-20 min followed by cooling at RT for 20 min.)

The optimal dilution for a specific application should be determined by the investigator.

Positive Control: Tonsil

Cellular Localization: Cell membrane and cytoplasm

Supplied As:

200µg/ml antibody purified from the ascites fluid by Protein G chromatography. Prepared in 10mM PBS, pH 7.4, with 0.2% BSA and 0.09% sodium azide. Also available without BSA and azide at 1mg/ml,

Prediluted antibody which is ready-to-use for staining of formalin-fixed, paraffin-embedded tissues.

Storage and Stability:

Ab with sodium azide is stable for 24 months when stored at 2-8°C. Antibody WITHOUT sodium azide is stable for 36 months when stored at below 0°C.

dianova GmbH Warburgstr. 45 ● 20354 Hamburg Telefon (040)45067-0 ● Telefax (040) 45067-490 ● www.dianova.de



DATA SHEET

CD1a Ab-5 (Clone O10)

Mouse Monoclonal Antibody

Catalog #DLN-14729, DLN-14730, or DLN-14733 (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) Cat. #DLN-14731 or DLN-14734 (0.1ml or 0.2ml at 1.0mg/ml) (Purified Ab without BSA and Azide) Catalog #DLN-14732 (7.0ml)

Suggested References:

- 1. Aractingi S, et al. J Clin Pathol. 50: 305-309 (1997).
- 2. Bobryshev Y V, et al. Pathol Res Pract. 192: 462-467 (1996).
- **3.** Mazal P R, et al Clin Neuropathol. 15: 87-91 (1996).

Limitations and Warranty:

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the actual price paid for the product. Dianova is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

Material Safety Data:

This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.

For Research Use Only