



Product Information

Catalog Number: DkxSh-003-DFITC

Product: Donkey anti-Sheep IgG (H&L) - Affinity Pure, FITC Conjugate

Physical State: Clear, fluorescent yellow liquid

Size: 2.0 mg

Antibody Concentration: 1.50 mg/ml (E 1% at 280 nm = 13.0) (concentration lot dependent)

Fluorophore: Fluorescein-5-isothiocyanate (FITC)

Amax = 494 nm; Emax = 518 nmFluorescein-5-isothiocyanate (FITC)

Suggested Dilution Range: 1:50 - 1:500 for most applications

The optimal working dilution should be determined by the investigator.

Buffer: 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Stabilizer: 1% (w/v) Bovine Serum Albumin (IgG-free, Protease-Free)

Preservative: 0.05% (w/v) Sodium Azide

Purification: Affinity purified using solid phase Sheep IgG

Purity: Affinity purified antibody is > 95% based on SDS-PAGE

Antibody Specificity: Based on IEP, this antibody reacts with:

· heavy (y) chains on sheep IgG

· light chains on all sheep immunoglobulins Based on IEP, no reactivity is observed to: · non-immunoglobulin sheep serum proteins

Reconstitution and Storage: Store undiluted liquid at 2-8°C. For storage in liquid form at -20°C, dilute with an equal volume

of glycerol (Molecular Grade, purity >99,5%) to prevent loss of activity. Product is stable for up to 1 year from date of receipt. Prepare working dilution prior to use and then discard. **Note:** adding glycerol reduces the stated protein concentration and dilution range by one-

nait.

Standard Laboratory Practices should be followed when handling this material. The chemical,

physical, and toxicological properties of this material have not been thoroughly investigated. The antibody contains 0.05% sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate

measures should be taken to avoid skin and eye contact, inhalation, and ingestion.

Note: This information sheet contains <u>general product information only</u>. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Warning: Bovine Serum Albumin (BSA) or dry milk is not suitable for blocking or diluting anti-bovine, anti-goat, anti-sheep or anti-horse IgG antibodies. BSA and dry milk may contain IgG, which react with these antibodies and may cause high background and/or reduce antibody titer. We recommend to use normal serum (5% v/v) from the same host species as the labeled secondary antibody.

Manufacturer:

ImmunoReagents, Inc. 6003 Chapel Hill Road, Suite 153 Raleigh, NC 27607, USA

Distribution and Support:

dianova GmbH Warburgstr. 45 20354 Hamburg www.dianova.com Technical Support: Phone: 040 45 06 70 Email: info@dianova.de Ordering Information:
Phone: 040 45 06 70
Fax: 040 45 06 74 90
Email: order@dianova.de