



Product Information

Catalog Number:	DkxSh-003-F
Product:	Donkey anti-Sheep IgG (H&L) - Affinity Pure, min x w/human, mouse, or rabbit IgG
Physical State:	Clear, colorless liquid, 0.2 µm filtered
Size:	2.0 mg
Antibody Concentration:	> 4.5 mg/ml (E 1% at 280 nm = 14.0) (concentration lot dependent)
Suggested Dilution Range:	10 - 20 μg/ml for most applications The optimal working dilution should be determined by the investigator.
Buffer:	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer:	none
Preservative:	0.05% (w/v) Sodium Azide
Purification:	Affinity purified using solid phase Sheep IgG
Purity:	> 90% based on SDS-PAGE Small amounts of intact IgG may be present.
Antibody Specificity:	Based on IEP, this antibody reacts with: · heavy (γ) chains on sheep IgG · light chains on all sheep immunoglobulins Based on IEP, no reactivity is observed to: · non-immunoglobulin sheep serum proteins · IgG from human, mouse or rabbit
Reconstitution and Storage:	Store at 2-8 °C. Product is stable for up to 1 year from date of receipt. Prepare working dilution prior to use and then discard.
Safety Information:	Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. The antibody contains 0.05% sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion.

Note: This information sheet contains <u>general product information only</u>. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Warning: Bovine Serum Albumin (BSA) or dry milk is not suitable for blocking or diluting anti-bovine, anti-goat, anti-sheep or anti-horse IgG antibodies. BSA and dry milk may contain IgG, which react with these antibodies and may cause high background and/or reduce antibody titer. We recommend to use normal serum (5% v/v) from the same host species as the labeled secondary antibody.

Manufacturer:

ImmunoReagents, Inc. 6003 Chapel Hill Road, Suite 153 Raleigh, NC 27607, USA

Distribution and Support:

dianova GmbH Warburgstr. 45 20354 Hamburg www.dianova.com Technical Support: Phone: 040 45 06 70 Email: info@dianova.de Ordering Information: Phone: 040 45 06 70 Fax: 040 45 06 74 90 Email: order@dianova.de