



## **Product Information**

| Catalog Number:             | GtxHu-004-GFITC   |
|-----------------------------|---|
| Product:                    | Goat anti-Human IgG Fc, F(ab)'2 Fragment - Affinity Pure, FITC Conjugate  |
| Physical State:             | Clear, fluorescent yellow liquid  |
| Size:                       | 0.5 mg  |
| Antibody Concentration:     | 1.50 mg/ml (E 1% at 280 nm = 13.0) (concentration lot dependent)  |
| Fluorophore:                | Fluorescein-5-isothiocyanate (FITC)<br>Amax = 494 nm; Emax = 518 nmFluorescein-5-isothiocyanate (FITC)  |
| Suggested Dilution Range:   | 1:50 - 1:500 for most applications<br>The optimal working dilution should be determined by the investigator.  |
| Buffer:                     | 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |
| Stabilizer:                 | 1% (w/v) Bovine Serum Albumin (IgG-free, Protease-Free)   |
| Preservative:               | 0.05% (w/v) Sodium Azide  |
| Purification:               | Affinity purified using solid phase Human IgG   |
| Purity:                     | > 90% based on SDS-PAGE Small amounts of intact IgG may be present.   |
| Antibody Specificity:       | Based on IEP, this antibody reacts with:<br>· heavy (γ) chains on human IgG<br>Based on IEP, no reactivity is observed to:<br>· non-immunoglobulin human serum immunoglobulins<br>· light chains on all human immunoglobulins   |
| Reconstitution and Storage: | Store undiluted liquid at 2-8°C. For storage in liquid form at -20°C, dilute with an equal volume of glycerol (Molecular Grade, purity >99,5%) to prevent loss of activity. Product is stable for up to 1 year from date of receipt. Prepare working dilution prior to use and then discard. <b>Note:</b> adding glycerol reduces the stated protein concentration and dilution range by one-half.  |
| Safety Information:         | Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. The antibody contains 0.05% sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. |

**Note:** This information sheet contains <u>general product information only</u>. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer:

ImmunoReagents, Inc. 6003 Chapel Hill Road, Suite 153 Raleigh, NC 27607, USA

**Distribution and Support:** 

dianova GmbH Warburgstr. 45 20354 Hamburg www.dianova.com Technical Support: Phone: 040 45 06 70 Email: info@dianova.de **Ordering Information:** Phone: 040 45 06 70 Fax: 040 45 06 74 90 Email: <u>order@dianova.de</u>