

## Product Information

|                                    |  |
|------------------------------------|--|
| <b>Catalog Number:</b>             | GtxHu-084-D650NHSX   |
| <b>Product:</b>                    | Goat anti-Human Kappa ( $\kappa$ ) Light Chain - affinity purified, DyLight®650 Conjugate  |
| <b>Physical State:</b>             | Lyophilized  |
| <b>Size:</b>                       | 0.5 mg   |
| <b>Antibody Concentration:</b>     | 1.0 mg/ml (E 1% at 280 nm = 13.0) (concentration lot dependent)  |
| <b>Fluorophore:</b>                | DyLight® 650<br>( $\text{Ex}$ = 652 nm; $\text{Em}$ = 672 nm)  |
| <b>Suggested Dilution Range:</b>   | 1:100 - 1:1000 for most applications<br>The optimal working dilution should be determined by the investigator.   |
| <b>Buffer:</b>                     | 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
| <b>Stabilizer:</b>                 | 1% (w/v) Bovine Serum Albumin (IgG-free, Protease-Free)  |
| <b>Preservative:</b>               | 0.05% (w/v) Sodium Azide   |
| <b>Purification:</b>               | Affinity purified using solid phase Human Kappa ( $\kappa$ ) Chain   |
| <b>Purity:</b>                     | Affinity purified antibody is > 95% based on SDS-PAGE  |
| <b>Antibody Specificity:</b>       | Based on IEP, this antibody reacts with: <ul style="list-style-type: none"><li>· kappa (<math>\kappa</math>) light chains on all human immunoglobulins</li></ul> Based on IEP, no reactivity is observed to: <ul style="list-style-type: none"><li>· non-immunoglobulin human serum immunoglobulins</li><li>· heavy chains on all human immunoglobulins</li><li>· lambda (<math>\lambda</math>) light chains on all human immunoglobulins</li></ul>  |
| <b>Reconstitution and Storage:</b> | Store freeze-dried powder at 2-8°C. Rehydrate with 0.55 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily. Product is stable for up to 4 weeks at 2-8°C after rehydration. For extended storage after rehydration, add an equal volume of glycerol (Molecular Grade, purity > 99.5%) for a final concentration of 50%, and store at -20°C in liquid form. <b>Note:</b> adding glycerol reduces the stated protein concentration and dilution range by one-half. Under long-term storage the rehydrated antibody is stable for one year from date of rehydration. The expiration date may be extended if test results are acceptable for the intended use. |
| <b>Safety Information:</b>         | Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. The antibody contains 0.05% sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion.  |

**Note:** This information sheet contains general product information only. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

DyLight® is a trademark of Thermo Fisher Scientific, Inc. and its subsidiaries.

**This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.**

### Manufacturer:

**ImmunoReagents, Inc.**  
6003 Chapel Hill Road, Suite 153  
Raleigh, NC 27607, USA

### Distribution and Support:

**dianova GmbH**  
Warburgstr. 45  
20354 Hamburg  
www.dianova.com

**Technical Support:**  
Phone: 040 45 06 70  
Email: [info@dianova.de](mailto:info@dianova.de)

**Ordering Information:**  
Phone: 040 45 06 70  
Fax: 040 45 06 74 90  
Email: [order@dianova.de](mailto:order@dianova.de)