

## Product Information

<b>Catalog Number:</b>	GtxMu-003-JHRPX
<b>Product:</b>	Goat anti-Mouse IgG (H&L), F(ab)'2 Fragment - Affinity Pure, HRP Conjugate, min x w/human IgG or serum proteins
<b>Physical State:</b>	Lyophilized
<b>Size:</b>	0.5 mg
<b>Antibody Concentration:</b>	1.0 mg/ml (E 1% at 280 nm = 13.2) (concentration lot dependent)
<b>Conjugation:</b>	Horseradish Peroxidase
<b>Suggested Dilution Range:</b>	1:500 - 1:5.000 for most applications The optimal working dilution should be determined by the investigator.
<b>Buffer:</b>	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Stabilizer:</b>	1% (w/v) Bovine Serum Albumin (IgG-free, Protease-Free)
<b>Preservative:</b>	0.1% (v/v) Kathon CG
<b>Purification:</b>	Affinity purified using solid phase Mouse IgG
<b>Purity:</b>	> 90% based on SDS-PAGE Small amounts of intact IgG may be present.
<b>Antibody Specificity:</b>	Based on IEP, this antibody reacts with: <ul style="list-style-type: none"><li>· heavy (γ) chains on mouse IgG</li><li>· light chains on all mouse immunoglobulins</li></ul> Based on IEP, no reactivity is observed to: <ul style="list-style-type: none"><li>· non-immunoglobulin mouse serum immunoglobulins</li><li>· human IgG or serum proteins</li></ul>
<b>Reconstitution and Storage:</b>	Store freeze-dried powder at 2-8 °C. Rehydrate with 0.55 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily. Product is stable for up to 4 weeks at 2-8°C after rehydration. For extended storage, aliquot and freeze at -70°C or below. Avoid repeated freezing and thawing. Alternatively, add an equal volume of glycerol (Molecular Grade, purity > 99,5%) and store at -20°C in liquid form. <b>Note:</b> adding glycerol reduces the stated protein concentration and dilution range by one-half. Under long-term storage the rehydrated antibody is stable for one year from date of rehydration. <b>Warning: Confirm that glycerol is non-contaminated and of high-quality (Molecular Grade, purity &gt; 99,5%) before adding to this product. Use of low quality and/ or contaminated glycerol may seriously inhibit peroxidase enzyme activity.</b>
<b>Safety Information:</b>	The product contains no hazardous constituents. Standard Laboratory Practices should be followed when handling this material.

**Note:** This information sheet contains general product information only. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

**This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.**

### Manufacturer:

**ImmunoReagents, Inc.**  
6003 Chapel Hill Road, Suite 153  
Raleigh, NC 27607, USA

### Distribution and Support:

**dianova GmbH**  
Warburgstr. 45  
20354 Hamburg  
www.dianova.com

**Technical Support:**  
Phone: 040 45 06 70  
Email: [info@dianova.de](mailto:info@dianova.de)

**Ordering Information:**  
Phone: 040 45 06 70  
Fax: 040 45 06 74 90  
Email: [order@dianova.de](mailto:order@dianova.de)