



## **Product Information**

**Catalog Number:** 101-155-165

Product: AMCA-AffiniPure Goat Anti-Bovine IgG (H+L) (minimal cross-reaction to Armenian Hamster,

Human, Mouse, and Rat Serum Proteins)

Physical State: Freeze-dried solid

**Size:** 1.5 mg

**Antibody Concentration:** ± 1.5 mg/ml (exact concentration lot dependent)

Fluorophore: 7-Amino-4-methylcoumarin-3-acetic acid-NHS ester

Amax = 350 nm; Emax = 450 nm

**Suggested Dilution Range:** 1:50 - 1:200 for most applications

**Buffer:** 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6

Stabilizer: 15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free)

**Preservative:** 0.05% Sodium Azide

Purity: The antibody was purified from antisera by immunoaffinity chromatography using antigens

coupled to agarose beads.

Antibody Specificity: Based on immunoelectrophoresis and/or ELISA, the antibody reacts with whole molecule

bovine IgG. It also reacts with the light chains of other bovine immunoglobulins. No antibody was detected against non immunoglobulin serum proteins. The antibody has been tested by ELISA and/or solid-phase adsorbed to ensure minimal cross-reaction with Armenian hamster, human, mouse, and rat serum proteins, but it may cross-react with immunoglobulins from other

species.

Reconstitution and Storage: Store freeze-dried powder at 2-8°C. When ready to use, rehydrate with 1.5 ml dH2O (exact

volume is lot specific) and centrifuge if not clear. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Prepare working dilution fresh each day. For extended storage after rehydration, aliquot and freeze at -70°C or below. Avoid repeated freezing and thawing. Alternatively, add an equal volume of glycerol (ACS grade or better) for a final concentration of 50%, and store at -20°C as a liquid. Note: adding glycerol reduces the stated protein

concentration and dilution range by one-half.

Expiration date:one year from date of rehydration. The expiration date may be extended if test

results are acceptable for the intended use.

**Safety Information:** Standard Laboratory Practices should be followed when handling this material. The chemical,

physical, and toxicological properties of this material have not been thoroughly investigated. The antibody contains 0.05% sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion.

**Note:** This information sheet contains general product information only. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Warning: Bovine serum albumin (BSA) and dry milk may contain IgG. Secondary antibodies directed against alpaca, bovine, goat, horse, or sheep react with bovine IgG. Therefore, use of BSA or milk to block or dilute these antibodies may significantly increase background and/or reduce antibody titer. For blocking, normal serum (5% v/v) from the host species of the labeled secondary antibody is recommended.

Manufacturer:

Jackson ImmunoResearch Laboratories, Inc. 872 West Baltimore Pike West Grove, PA, USA 19390

**Distribution and Support:** 

dianova GmbH Warburgstraße 45 20354 Hamburg www.dianova.com Technical Support: Phone: 040 45 06 70 Email: info@dianova.de Ordering Information:
Phone: 040 45 06 70
Fax: 040 45 06 74 90
Email: order@dianova.de