

## Product Information

|                                    |  |
|------------------------------------|--|
| <b>Catalog Number:</b>             | 109-476-170  |
| <b>Product:</b>                    | DyLight™ 405 AffiniPure F(ab') <sub>2</sub> Fragment Goat Anti-Human IgG, Fcγ fragment specific (minimal cross-reaction to Bovine, Mouse, and Rabbit Serum Proteins)   |
| <b>Physical State:</b>             | Freeze-dried solid   |
| <b>Size:</b>                       | 0.75 mg  |
| <b>Antibody Concentration:</b>     | 1.4 - 1.5 mg/ml (exact concentration lot dependent)  |
| <b>Fluorophore:</b>                | DyLight™405-TFP ester<br>Amax=400; Emax=420  |
| <b>Suggested Dilution Range:</b>   | 1:100 - 1:800 for most applications  |
| <b>Buffer:</b>                     | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6   |
| <b>Stabilizer:</b>                 | 15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free)  |
| <b>Preservative:</b>               | 0.05% Sodium Azide   |
| <b>Purity:</b>                     | The antibody was purified from antisera by a combination of pepsin digestion and immunoaffinity chromatography using antigens coupled to agarose beads. Fc fragments and whole IgG molecules have been removed.  |
| <b>Antibody Specificity:</b>       | Based on immunoelectrophoresis and/or ELISA, the antibody reacts with the Fc portion of human IgG heavy chain but not with the Fab portion of human IgG. No antibody was detected against human IgM or IgA, or against non-immunoglobulin serum proteins. The antibody has been tested by ELISA and/or solid-phase adsorbed to ensure minimal cross-reaction with bovine, mouse and rabbit serum proteins, but it may cross-react with immunoglobulins from other species.   |
| <b>Reconstitution and Storage:</b> | Store freeze-dried powder at 2-8°C. When ready to use, rehydrate with 0.75 ml dH <sub>2</sub> O (exact volume is lot specific) and centrifuge if not clear. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Prepare working dilution fresh each day. For extended storage after rehydration, aliquot and freeze at -70°C or below. Avoid repeated freezing and thawing. Alternatively, add an equal volume of glycerol (ACS grade or better) for a final concentration of 50%, and store at -20°C as a liquid. Note: adding glycerol reduces the stated protein concentration and dilution range by one-half.<br>Expiration date: one year from date of rehydration. The expiration date may be extended if test results are acceptable for the intended use. |
| <b>Safety Information:</b>         | Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. The antibody contains 0.05% sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion.  |

**Note:** This information sheet contains general product information only. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

**This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.**

### Manufacturer:

**Jackson ImmunoResearch Laboratories, Inc.**  
872 West Baltimore Pike  
West Grove, PA, USA 19390

### Distribution and Support:

**dianova GmbH**  
Warburgstraße 45  
20354 Hamburg  
www.dianova.com

**Technical Support:**  
Phone: 040 45 06 70  
Email: [info@dianova.de](mailto:info@dianova.de)

**Ordering Information:**  
Phone: 040 45 06 70  
Fax: 040 45 06 74 90  
Email: [order@dianova.de](mailto:order@dianova.de)