

Product Information

Catalog Number:	205-002-176
Product:	IgG Fraction Monoclonal Mouse Anti-Goat IgG, Light Chain Specific (minimal cross-reaction to Horse, Human, Mouse, Rabbit, and Rat Immunoglobulins) Clone 20B07-02A09 (IgG1/k)
Physical State:	Sterile-filtered liquid
Size:	1.0 mg
IgG Concentration:	1.2 - 1.3 mg/ml (exact concentration lot dependent)
Conjugation:	unconjugated
Suggested Working Concentration:	10-20 ug/ml
Buffer:	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6
Stabilizer:	None, BSA-free
Preservative:	None
Purity:	The antibody was purified from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.
Antibody Specificity:	Based on antigen-binding assay, Western blotting, and/or ELISA, the antibody reacts with the light chains on goat IgG and with those common to other goat immunoglobulins. The antibody does not react with the heavy chain of goat IgG. The antibody has been tested by ELISA to ensure minimal cross-reaction with horse, human, mouse, rabbit, and rat immunoglobulins, but it may cross-react with immunoglobulins from other species.
Storage:	Store at 2-8°C under sterile conditions. Prepare working dilution fresh each day. Expiration date: one year from date of receipt. The expiration date may be extended if test results are acceptable for the intended use.
Safety Information:	This product contains no substances considered to be hazardous to health. Although the product is classified as non-hazardous, Standard Laboratory Practices should be followed when handling this material.

Note: This information sheet contains general product information only. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Warning: Bovine serum albumin (BSA) and dry milk may contain IgG. Secondary antibodies directed against alpaca, bovine, goat, horse, or sheep react with bovine IgG. Therefore, use of BSA or milk to block or dilute these antibodies may significantly increase background and/or reduce antibody titer. For blocking, normal serum (5% v/v) from the host species of the labeled secondary antibody is recommended.

Manufacturer:

Jackson ImmunoResearch Laboratories, Inc.
872 West Baltimore Pike
West Grove, PA, USA 19390

Distribution and Support:

dianova GmbH
Warburgstraße 45
20354 Hamburg
www.dianova.com

Technical Support:
Phone: 040 45 06 70
Email: info@dianova.de

Ordering Information:
Phone: 040 45 06 70
Fax: 040 45 06 74 90
Email: order@dianova.de