

## Product Information

<b>Catalog Number:</b>	305-006-003
<b>Product:</b>	AffiniPure F(ab') <sub>2</sub> Fragment Rabbit Anti-Goat IgG (H+L)
<b>Physical State:</b>	Sterile-filtered liquid
<b>Size:</b>	1.0 mg
<b>Antibody Concentration:</b>	1.2 - 1.3 mg/ml (exact concentration lot dependent)
<b>Conjugation:</b>	unconjugated
<b>Suggested Working Concentration:</b>	10-20 ug/ml
<b>Buffer:</b>	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6
<b>Stabilizer:</b>	None, BSA-free
<b>Preservative:</b>	None
<b>Purity:</b>	The antibody was purified from antisera by a combination of pepsin digestion and immunoaffinity chromatography using antigens coupled to agarose beads. Fc fragments and whole IgG molecules have been removed.
<b>Antibody Specificity:</b>	Based on immunoelectrophoresis and/or ELISA, the antibody reacts with whole molecule goat IgG. It also reacts with the light chains of other goat immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. The antibody may cross-react with immunoglobulins from other species.
<b>Storage:</b>	Store at 2-8°C under sterile conditions. Prepare working dilution fresh each day. Expiration date: one year from date of receipt. The expiration date may be extended if test results are acceptable for the intended use.
<b>Safety Information:</b>	This product contains no substances considered to be hazardous to health. Although the product is classified as non-hazardous, Standard Laboratory Practices should be followed when handling this material.

**Note:** This information sheet contains general product information only. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

**This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.**

**Warning:** Bovine serum albumin (BSA) and dry milk may contain IgG. Secondary antibodies directed against alpaca, bovine, goat, horse, or sheep react with bovine IgG. Therefore, use of BSA or milk to block or dilute these antibodies may significantly increase background and/or reduce antibody titer. For blocking, normal serum (5% v/v) from the host species of the labeled secondary antibody is recommended.

### Manufacturer:

**Jackson ImmunoResearch Laboratories, Inc.**  
872 West Baltimore Pike  
West Grove, PA, USA 19390

### Distribution and Support:

**dianova GmbH**  
Warburgstraße 45  
20354 Hamburg  
www.dianova.com

**Technical Support:**  
Phone: 040 45 06 70  
Email: [info@dianova.de](mailto:info@dianova.de)

**Ordering Information:**  
Phone: 040 45 06 70  
Fax: 040 45 06 74 90  
Email: [order@dianova.de](mailto:order@dianova.de)