



Product Information

Catalog Number:	305-545-047
Product:	Alexa Fluor® 488-AffiniPure Rabbit Anti-Goat IgG, F(ab')2 Fragment Specific (minimal cross- reaction to Human Serum Proteins)
Physical State:	Freeze-dried solid
Size:	1.0 mg
Antibody Concentration:	1.4 - 1.5 mg/ml (exact concentration lot dependent)
Fluorophore:	Alexa Fluor®488 carboxylic acid Amax = 493 nm; Emax = 519 nm
Suggested Dilution Range:	1:100 - 1:800 for most applications
Buffer:	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6
Stabilizer:	15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free)
Preservative:	0.05% Sodium Azide
Purity:	The antibody was purified from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.
Antibody Specificity:	Based on immunoelectrophoresis and/or ELISA, the antibody reacts with the F(ab')2/Fab
Antibody Specificity.	portion of goat IgG. It also reacts with the light chains of other goat immunoglobulins. No antibody was detected against the Fc portion of goat IgG or against non-immunoglobulin serum proteins. The antibody has been tested by ELISA and/or solid phase adsorbed to ensure minimal cross-reaction with human serum proteins, but it may cross-react with immunoglobulins from other species.
Reconstitution and Storage:	portion of goat IgG. It also reacts with the light chains of other goat immunoglobulins. No antibody was detected against the Fc portion of goat IgG or against non-immunoglobulin serum proteins. The antibody has been tested by ELISA and/or solid phase adsorbed to ensure minimal cross-reaction with human serum proteins, but it may cross-react with

Note: This information sheet contains general product information only. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Warning: Bovine serum albumin (BSA) and dry milk may contain IgG. Secondary antibodies directed against alpaca, bovine, goat, horse, or sheep react with bovine IgG. Therefore, use of BSA or milk to block or dilute these antibodies may significantly increase background and/or reduce antibody titer. For blocking, normal serum (5% v/v) from the host species of the labeled secondary antibody is recommended.

Manufacturer:

Jackson ImmunoResearch Laboratories, Inc. 872 West Baltimore Pike West Grove, PA, USA 19390

Distribution and Support:

dianova GmbH Warburgstraße 45 20354 Hamburg www.dianova.com Technical Support: Phone: 040 45 06 70 Email: info@dianova.de Ordering Information: Phone: 040 45 06 70 Fax: 040 45 06 74 90 Email: order@dianova.de