



## **Product Information**

Catalog Number: 705-035-147

**Product:** Peroxidase-AffiniPure Donkey Anti-Goat IgG (H+L) (minimal cross-reaction to Chicken, Guinea

Pig, Syrian Hamster, Horse, Human, Mouse, Rabbit, and Rat Serum Proteins)

**Physical State:** Freeze-dried solid

Size: 0.5 ml

**Antibody Concentration:** ± 0.8 mg/ml (exact concentration lot dependent)

Conjugation: Horseradish Peroxidase (HRP)

1:500 - 1:5.000 for immunohisto/cytochemistry Suggested Dilution Range:

1:5.000 - 1:100.000 for ELISA and Western blotting with chromogenic substrates

1:10.000 - 1:200.000 for Western blotting with ECL substrates

**Buffer:** 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6

Stabilizer: 15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free)

Preservative: None (Warning: Use of sodium azide as a preservative will substantially inhibit the enzyme

activity of horseradish peroxidase.)

**Purity:** The antibody was purified from antisera by immunoaffinity chromatography using antigens

coupled to agarose beads.

**Antibody Specificity:** Based on immunoelectrophoresis and/or ELISA, the antibody reacts with whole molecule goat

IgG. It also reacts with the light chains of other goat immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. The antibody has been tested by ELISA and/or solid-phase adsorbed to ensure minimal cross-reaction with chicken, guinea pig, Syrian hamster, horse, human, mouse, rabbit, and rat serum proteins, but it may cross-react with

immunoglobulins from other species.

Reconstitution and Storage: Store freeze-dried powder at 2-8°C. When ready to use, rehydrate with 0.5 ml dH2O and

centrifuge if not clear. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Prepare working dilution fresh each day. For extended storage after rehydration, aliquot and freeze at -70°C or below. Avoid repeated freezing and thawing. Alternatively, add an equal volume of glycerol (ACS grade or better) for a final concentration of 50%, and store at -20°C as a liquid. Note: adding glycerol reduces the stated protein concentration and dilution range by one-half.

Expiration date:one year from date of rehydration. The expiration date may be extended if test

results are acceptable for the intended use.

(Warning: Confirm that glycerol is ACS grade or better before adding to this product. Lower

grades of glycerol may seriously inhibit peroxidase enzyme activity.)

Safety Information: This product contains no substances considered to be hazardous to health. Although the

product is classified as non-hazardous, Standard Laboratory Practices should be followed

when handling this material.

Note: This information sheet contains general product information only. For detailed lot specific information consult the vial label and the data sheet supplied by the manufacturer upon delivery of the product.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Warning: Bovine serum albumin (BSA) and dry milk may contain IgG. Secondary antibodies directed against alpaca, bovine, goat, horse, or sheep react with bovine IgG. Therefore, use of BSA or milk to block or dilute these antibodies may significantly increase background and/or reduce antibody titer. For blocking, normal serum (5% v/v) from the host species of the labeled secondary antibody is recommended.

Manufacturer:

Jackson ImmunoResearch Laboratories, Inc. 872 West Baltimore Pike West Grove, PA, USA 19390

**Distribution and Support:** 

dianova GmbH Warburgstraße 45 20354 Hamburg www.dianova.com **Technical Support:** Phone: 040 45 06 70 Email: info@dianova.de

Ordering Information: Phone: 040 45 06 70 Fax: 040 45 06 74 90 Email: order@dianova.de





## Manufacturer:

Jackson ImmunoResearch Laboratories, Inc. 872 West Baltimore Pike West Grove, PA, USA 19390

**Distribution and Support:** 

dianova GmbH Warburgstr. 45 20354 Hamburg www.dianova.com Technical Support: Phone: 040 45 06 70 Email: info@dianova.de Ordering Information: Phone: 040 45 06 70 Fax: 040 45 06 74 90 Email: order@dianova.de