

Beratung und Vertrieb

Fon + 49 (0) 40 45 067 0

dianova

Fax + 49 (0) 40 450 67 490

info@dianova.de

Warburgstr. 45 • 20354 Hamburg

www.dianova.de

Product: Rabbit Monoclonal Antibody
Biotin Anti-Mouse IgG Rabbit Monoclonal Antibody,
Clone RM104

Catalog No.: 31-1001-02

Lot No.:

Clone RM104

Specificity This antibody reacts to the Fc region of all mouse IgG subclasses. No cross reactivity with mouse IgM, IgA, IgE, human IgG, and rat IgG. It may cross react to goat IgG.

The Fc region of RM104 has been engineered to eliminate Fc receptor binding.

Application: ELISA, Flow Cytometry, Immunocytochemistry, Immunohistochemistry, Immunoprecipitation, Western Blot (nonreduced).

Immunogen: Mouse IgG

Purity: Protein A affinity purified from an animal origin-free and protein-free culture supernatant

Size: 50 µg

Concentration: 1.0 mg/mL


Buffer: 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide

Usage: ELISA: 0.005µg/mL – 0.2µg/mL;
ICC, IHC: 0.5µg/mL – 2µg/mL;
WB: 0.1µg/mL – 0.5µg/mL


Storage and Stability: Stable for 1 Year at -20.0°C from date of receipt.

Country of Origin: U.S.A.


Intended Use: For Research Use Only Not for Diagnostic or Therapeutic Use



ELISA of mouse immunoglobulins shows RM104 reacts to the Fc region of mouse IgG1, IgG2a, IgG2b, and IgG3; no cross reactivity with IgM, IgA, IgE, human or rat IgG. The plate was coated with 50 ng/well of different immunoglobulins. 200 ng/mL, 50 ng/mL, or 10 ng/mL of RM104 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.



A titer ELISA of mouse IgG. The plate was coated with different amounts of mouse IgG. A serial dilution of RM104 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.



Western blot of nonreduced(-) and reduced(+) mouse immunoglobulins (20ng/lane), using 0.2µg/mL of RevMAB clone RM104. This antibody reacts to nonreduced mouse IgG1, IgG2a, IgG2b, and IgG3. It showed no cross reactivity with IgM, IgA, or IgE.