

RevMAb Biosciences USA, Inc. 870 Dubuque Ave, South San Francisco, CA 94080, USA

Certificate of Analysis

Product: Rabbit Monoclonal Antibody

Anti-Mouse Immunoglobulin Kappa Light Chain Rabbit

Monoclonal Antibody, Clone RM103

Catalog No.: 31-1007-00

Lot No.:

Clone RM103

Specificity This antibody reacts to the kappa light chain of mouse

immunoglobulins. No cross reactivity with the lamda

light chain, human IgG, rat IgG, or goat IgG.

The Fc region of RM103 has been engineered to

eliminate Fc receptor binding.

Application: ELISA, Flow Cytometry, Immunocytochemistry,

Immunohistochemistry, Immunoprecipitation, Western

Blot (nonreduced).

Immunogen: Mouse IgG

Purity: Protein A affinity purified from an animal origin–free

and protein-free culture supernatant

Size: 100 μg

Concentration: 1.0 mg/mL

Buffer: 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide

Usage: ELISA: 0.005ug/mL – 0.2ug/mL;

ICC, IHC: 0.5ug/mL - 2ug/mL; WB: 0.1ug/mL - 0.5ug/mL

Storage and

Stable for 1 Year at -20.0°C from date of receipt.

Stability:

Country of Origin: U.S.A.

Warburgstr. 45 • 20354 Hamburg


Intended Use: For Research Use Only Not for Diagnostic or

Therapeutic Use


Beratung und Vertrieb Fon + 49 (0) 40 45 067 0

Fax + 49 (0) 40 450 67 490


info@dianova.de www.dianova.de



ELISA of mouse immunoglobulins shows RM103 reacts to the kappa light chain of mouse immunoglobulins. No cross reactivity with the lamda light chain, human lgG, rat lgG, or goat lgG. The plate was coated with 50 ng/well of different immunoglobulins. 200 ng/mL, 50 ng/mL, or 10 ng/mL of RM103 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit lgG as the secondary antibody.



A titer ELISA of mouse $IgG1\kappa$. The plate was coated with different amounts of mouse $IgG1\kappa$. A serial dilution of RM103 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.



Western blot of nonreduced(-) and reduced(+) mouse $IgG1\kappa$ and $IgG1\lambda$ (20ng/lane), using 0.2ug/mL of RevMAb clone RM103. This antibody reacts to nonreduced $IgG1\kappa$ (~150 kDa), and slightly reacts to reduced κ light chain (~25 kDa).