

RevMAb Biosciences USA, Inc. 870 Dubuque Ave, South San Francisco, CA 94080, USA

Beratung und Vertrieb

Fon + 49 (0) 40 45 067 0

diemovo

Fax + 49 (0) 40 450 67 490

Warburgstr. 45 • 20354 Hamburg

info@dianova.de www.dianova.de

Product:

Rabbit Monoclonal Antibody

Biotin Anti-Human IgG1 Rabbit Monoclonal Antibody,

Clone RM117

Catalog No.:

31-1019-02

Lot No.:

Clone

RM117

Specificity

This antibody reacts to the heavy chain of human IgG1.

No cross reactivity with human IgG2, IgG3, IgG4, IgM,

IgA, IgD, IgE, mouse IgG, rat IgG, or goat IgG.

Application:

ELISA, Immunohistochemistry, Immunocytochemistry,

Flow Cytometry.

Immunogen:

Peptide corresponding to the hinge region of Human

lgG1

**Purity:** 

Protein A affinity purified from an animal origin–free

culture supernatant

**Size:** 50 μg

Concentration: 1.0 mg/mL

**Buffer:** 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide

**Usage:** ELISA: 50ng/well – 200ng/well (for Capture);

0.05ug/mL - 0.2ug/mL (for Detection);

IHC, ICC: 0.5ug/mL - 2ug/mL;

Storage and


Stability:

Stable for 1 Year at -20.0°C from date of receipt.


Country of Origin: U.S.A.

Intended Use: For Research Use Only Not for Diagnostic or


**Therapeutic Use** 



Sandwich ELISA using RM117 as the capture antibody, and Biotinylated anti-human light chains  $(\kappa+\lambda)$  antibody RM129 as the detection antibody, followed by an AP conjugated streptavidin.



ELISA of human immunoglobulins shows RM117 only reacts to human IgG1. No cross reactivity with Human IgG2, IgG3, IgG4, IgE, IgD, IgA, mouse IgG, rat IgG, or goat IgG. The plate was coated with 50 ng/well of different immunoglobulins. 200 ng/mL, 50 ng/mL, or 10 ng/mL of RM117 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.



A titer ELISA using RM117. The plate was coated with different amounts of human IgG1. A serial dilution of RM117 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.