

RevMAb Biosciences USA, Inc. 830 Dubuque Ave, South San Francisco, CA 94080, USA

Certificate of Analysis

Product: Rabbit Monoclonal Antibody

Human IgD Matched Antibody Pair

Catalog No.: 31-1025-MK

Lot No.:

Specificity This antibody pair detects only human IgD. It does not

react to monkey (Cyno or Rhesus) IgD, mouse IgD, rat

IgD, or goat IgD.

Application: Sandwich ELISA, or other Sandwich Assays.

Components: 1) Capture Antibody: Anti-Human IgD Rabbit

Monoclonal antibody, clone RM123, 100 µg at 1.0 mg/mL in 50% Glycerol/PBS with 1% BSA and

0.09% sodium azide;

 Detection Antibody: Biotin Anti-Human Ig Light Chain Rabbit Monoclonal Antibody, Clone RM129,

25 μg at 1.0 mg/mL in 50% Glycerol/PBS with 1%

BSA and 0.09% sodium azide;

3) 10X Sample Diluent: Added to the tested sample to

reduce non-specific background and noise.

Usage: ELISA: Capture Antibody 50ng/well – 200ng/well;

Detection Antibody 0.1ug/mL - 0.5ug/mL;

Storage and Stability: Components (1) (2) Stable for 1 Year at -20.0°C from

date of receipt.

Component (3) Store at 2-8°C

Country of Origin: U.S.A.

Intended Use: For Research Use Only Not for Diagnostic or

Therapeutic Use

Beratung und Vertrieb

Warburgstr. 45 • 20354 Hamburg

Fon + 49 (0) 40 45 067 0 Fax + 49 (0) 40 450 67 490 info@dianova.de www.dianova.de




Fig 1. Sandwich ELISA using RM123 as the capture antibody (100ng/well), and Biotinylated anti-human light chains (k+l) antibody RM129 (0.2ug/mL) as the detection antibody, followed by an alkaline phosphatase conjugated streptavidin.




Fig 2. Sandwich ELISA, using RevMAb human IgD matched antibody pair (capture: RM123; detection: Biotin-RM129), shows species reactivity to human only, and shows no cross-reactivity to monkey (Cyno or Rhesus), mouse IgD, rat IgD, or goat IgD.




Fig 3. Sandwich ELISA, using RevMAb human IgD matched antibody pair, shows no cross-reactivity with Human IgG, IgE, IgA, or IgM. The plate was coated with human IgD capture antibody (50ng/well). Different immunoglobulin samples (0.2 ug/mL) were added and biotinylated RM129 (0.2ug/mL) was used as detection antibody, followed by an alkaline phosphatase conjugated streptavidin