

RevMAb Biosciences USA, Inc. 870 Dubuque Ave, South San Francisco, CA 94080, USA

Certificate of Analysis

Product: Rabbit Monoclonal Antibody

Anti-Human IgA1 Rabbit Monoclonal Antibody,

Clone RM124

Catalog No.: 31-1026-00

Lot No.:

Clone RM124

Specificity This antibody reacts to human IgA1, and very slightly

cross reacts with IgA2. No cross reactivity with human

IgG, IgM, IgD, or IgE.

Application: ELISA, Immunohistochemistry, Immunocytochemistry,

Flow Cytometry.

Immunogen: Human IgA

Purity: Protein A affinity purified from an animal origin-free

culture supernatant

Size: 100 µg

Concentration: 1.0 mg/mL

Buffer: 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide

ELISA: 50ng/well - 200ng/well (for Capture); Usage:

0.05ug/mL - 0.2ug/mL (for Detection);

ICC: 0.5ug/mL-2ug/mL;

IHC: 0.1ug/mL - 1ug/mL

Storage and

Stable for 1 Year at -20.0°C from date of receipt.

Stability:

Country of Origin: U.S.A.

Intended Use: For Research Use Only Not for Diagnostic or

Therapeutic Use


Beratung und Vertrieb


Fon + 49 (0) 40 45 067 0

Fax + 49 (0) 40 450 67 490


info@dianova.de

Warburgstr. 45 • 20354 Hamburg www.dianova.de







Sandwich ELISA using RM124 as the capture antibody (100ng/well), and Biotinylated antihuman light chains ($\kappa+\lambda$) antibody RM129 as the detection antibody, followed by an alkaline phosphatase conjugated streptavidin.



ELISA of human immunoglobulins shows RM124 only reacts to human IgA1. The clone very slightly cross reacts with IgA2. No cross reactivity with human IgG, IgM, IgD, or IgE. The plate was coated with 50 ng/well of different immunoglobulins. 200 ng/mL, 50ng/mL, or 10 ng/mL of RM124 was used as the primary antibody. An alkaline phosphatase conjugated antirabbit IgG as the secondary antibody.



Immunohistochemistry of Human Tonsil Tissue using Anti-Human IgA1 antibody RM124.



A titer ELISA using RM124. The plate was coated with different amounts of human IgA1. A serial dilution of RM124 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.