

RevMAb Biosciences USA, Inc. 870 Dubuque Ave, South San Francisco, CA 94080, USA

Beratung und Vertrieb

Fon + 49 (0) 40 45 067 0

Fax + 49 (0) 40 450 67 490

info@dianova.de

Warburgstr. 45 • 20354 Hamburg www.dianova.de

**Product:** Rabbit Monoclonal Antibody

Biotin Anti-Human IgA1 Rabbit Monoclonal Antibody,

Clone RM124

**Catalog No.:** 31-1026-02

Lot No.:

Clone RM124

**Specificity** This antibody reacts to human IgA1, and very slightly

cross reacts with IgA2. No cross reactivity with human

IgG, IgM, IgD, or IgE.

**Application:** ELISA, Immunohistochemistry, Immunocytochemistry,

Flow Cytometry.

Immunogen: Human IgA

**Purity:** Protein A affinity purified from an animal origin–free

culture supernatant

**Size:** 50 μg

**Concentration:** 1.0 mg/mL

**Buffer:** 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide

**Usage:** ELISA: 50ng/well – 200ng/well (for Capture);

0.05ug/mL - 0.2ug/mL (for Detection);

ICC: 0.5ug/mL-2ug/mL; IHC: 0.1ug/mL – 1ug/mL

Storage and


Stable for 1 Year at -20.0°C from date of receipt.


Stability:

Country of Origin: U.S.A.


Intended Use: For Research Use Only Not for Diagnostic or

Therapeutic Use







Sandwich ELISA using RM124 as the capture antibody (100ng/well), and Biotinylated antihuman light chains ( $\kappa+\lambda$ ) antibody RM129 as the detection antibody, followed by an alkaline phosphatase conjugated streptavidin.



ELISA of human immunoglobulins shows RM124 only reacts to human IgA1. The clone very slightly cross reacts with IgA2. No cross reactivity with human IgG, IgM, IgD, or IgE. The plate was coated with 50 ng/well of different immunoglobulins. 200 ng/mL, 50ng/mL, or 10 ng/mL of RM124 was used as the primary antibody. An alkaline phosphatase conjugated antirabbit IgG as the secondary antibody.



Immunohistochemistry of Human Tonsil Tissue using Anti-Human IgA1 antibody RM124.



A titer ELISA using RM124. The plate was coated with different amounts of human IgA1. A serial dilution of RM124 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.