



# Rabbit IgG anti-Hamster generally IgG (F(ab')2)-FITC, MinX none

# **General Information**

Catalog Number: SE	C-182546	Physical State:	Lyophilized
Quantity:	20 mg	Reconstitution Volume:	2.0 mL
		<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)
		Shipping Conditions:	at room temperature
		Product Expiration	Expiration date is one (1) year from date of opening.
Antibody Host / Forn	nat		
Host Species: Ra	bbit	Clonality / Isotype / Clone: polyclonal Ig	
Antibody Format: Igo	G whole molecule	Concentration:	
		Conjugation:	FITC (Fluoresceinisothiocyanat)
		Maximum Excitation / Emission: 492 nm / 520 nm	
Specificity			
Target Species:	Hamster generally	Immunogen:	Hamster IgG whole molecule
Antibody Specificity: IgG (F(ab')2 Fragment)		This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophores resulted in a single precipitin arc against anti-fluorescein, anti-Rabbit Serum, Golden Syrian Hamster IgG, Golden Syrian Hamster IgG F(ab')2 and Golden Syrian Hamster Serum. No reaction was observed against Golden Syrian Hamster IgG F(c).	
	Fragment)	fractionation and ion exc dialysis against the buffer resulted in a single precip Serum, Golden Syrian Han and Golden Syrian Hamst	hange chromatography followed by extensive stated above. Assay by immunoelectrophores itin arc against anti-fluorescein, anti-Rabbit mster IgG, Golden Syrian Hamster IgG F(ab')2 er Serum. No reaction was observed against
	Fragment)	fractionation and ion exc dialysis against the buffer resulted in a single precip Serum, Golden Syrian Han and Golden Syrian Hamst	hange chromatography followed by extensive stated above. Assay by immunoelectrophores itin arc against anti-fluorescein, anti-Rabbit mster IgG, Golden Syrian Hamster IgG F(ab')2 er Serum. No reaction was observed against G F(c).
Formulation, Transpo		fractionation and ion excl dialysis against the buffer resulted in a single precip Serum, Golden Syrian Han and Golden Syrian Hamster Ig	hange chromatography followed by extensive stated above. Assay by immunoelectrophores itin arc against anti-fluorescein, anti-Rabbit mster IgG, Golden Syrian Hamster IgG F(ab')2 er Serum. No reaction was observed against G F(c).
Storage Buffer: 0.02 N Phosphate, 0.15 M Sod	ort and Storage И Potassium	fractionation and ion excl dialysis against the buffer resulted in a single precip Serum, Golden Syrian Hamst Golden Syrian Hamster Ig <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge proc	hange chromatography followed by extensive stated above. Assay by immunoelectrophores itin arc against anti-fluorescein, anti-Rabbit mster IgG, Golden Syrian Hamster IgG F(ab')2 er Serum. No reaction was observed against G F(c). : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and luct if not completely clear after standing at
Storage Buffer: 0.02 M Phosphate, 0.15 M Sodi Stabilizer: None	ort and Storage И Potassium	fractionation and ion excl dialysis against the buffer resulted in a single precip Serum, Golden Syrian Ham and Golden Syrian Hamster Golden Syrian Hamster Ig <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge proc room temperature. This	hange chromatography followed by extensive stated above. Assay by immunoelectrophores itin arc against anti-fluorescein, anti-Rabbit mster IgG, Golden Syrian Hamster IgG F(ab')2 er Serum. No reaction was observed against G F(c). : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and luct if not completely clear after standing at
Storage Buffer: 0.02 N Phosphate, 0.15 M Sod	ort and Storage И Potassium	fractionation and ion excl dialysis against the buffer resulted in a single precip Serum, Golden Syrian Ham and Golden Syrian Hamster Golden Syrian Hamster Ig <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge proc room temperature. This	hange chromatography followed by extensive stated above. Assay by immunoelectrophores itin arc against anti-fluorescein, anti-Rabbit mster IgG, Golden Syrian Hamster IgG F(ab')2 er Serum. No reaction was observed against G F(c). : none restoration. For extended storage aliquot D° C or below. Avoid cycles of freezing and luct if not completely clear after standing at product is stable for several weeks at 4° C as an

Application: FLISA, Flow Cytometry, Immunofluorescence

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use. Management System Manufacturer BIOZOL ISO 9001:2015 Phone +49 (89) 3799 666 6 www.dianova.com Diagnostica Vertrieb GmbH Fax +49 (89) 3799 666 99 TÜVRheinland Your Secondary Antibody Portal Leipziger Straße 4 E-mail info@biozol.de ZERTIFIZIERT www.tuv.com ID 9000019771 Powered by BIOZOL 85386 Eching www.biozol.de



#### Product Datasheet / Instruction for use



SEC-182546 2023-02-23 2 / 2

Dilution: FLISA 1:10.000 - 1:50.000, Flow cytometry 1:500 - 1:2,500, Immunofluorescence 1:1,000 - 1:5,000

### **Application Note:**

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

# **Background Information**

Anti-Golden Syrian Hamster IgG F(ab')2 Fluorescein Antibody generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. F(ab')2 molecules lack the Fc portion of IgG and therefore receptors that bind Golden Syrian Hamster IgG F(c) will not bind Golden Syrian Hamster IgG F(ab')2 molecules. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

# Safety Information

The product contains no hazardous constituents. Standard Laboratory Practices should be followed when handling this material.

### Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL **Diagnostica Vertrieb GmbH** Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT www.tuv.com ID 9000019771

ISO 9001:2015

Management

System