

## Goat IgG anti-Mouse IgG (Fc)-HRPO, MinX none

### General Information

|                                   |  |
|-----------------------------------|--|
| <b>Catalog Number:</b> SEC-182588 | <b>Physical State:</b> Lyophilized   |
| <b>Quantity:</b> 20 mg            | <b>Reconstitution Volume:</b> 2.0 mL   |
|                                   | <b>Reconstitution Buffer:</b> Restore with deionized water (or equivalent)       |
|                                   | <b>Shipping Conditions:</b> at room temperature                                  |
|                                   | <b>Product Expiration:</b> Expiration date is one (1) year from date of opening. |

### Antibody Host / Format

|  |   |
|--|---|
| <b>Host Species:</b> Goat                  | <b>Clonality / Isotype / Clone:</b> polyclonal Ig           |
| <b>Antibody Format:</b> IgG whole molecule | <b>Concentration:</b> 10.0 mg/mL by UV absorbance at 280 nm |
|  | <b>Conjugation:</b> Horseradish Peroxidase (HRP)            |
|  | <b>Maximum Excitation / Emission:</b> /                     |

### Specificity

|  |   |
|--|---|
| <b>Target Species:</b> Mouse                   | <b>Immunogen:</b> Anti-Mouse IgG F(c) was produced by repeated immunization with mouse IgG F(c) fragment in goat.   |
| <b>Antibody Specificity:</b> IgG (Fc Fragment) | This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG F(ab). |
|  | <b>Minimal Cross Reactivity:</b> none   |

### Formulation, Transport and Storage

|   |   |
|---|---|
| <b>Storage Buffer:</b> 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2            | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| <b>Stabilizer:</b> 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |   |
| <b>Preservatives:</b> 0.01% (w/v) Gentamicin Sulfate                                      |   |
|   | <b>Storage Temperature:</b> 4-8°C   |

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

#### Manufacturer

**BIOZOL**  
Diagnostica Vertrieb GmbH  
Leipziger Straße 4  
85386 Eching

Phone +49 (89) 3799 666 6  
Fax +49 (89) 3799 666 99  
E-mail info@biozol.de  
www.biozol.de

[www.dianova.com](http://www.dianova.com)

Your Secondary Antibody Portal  
Powered by BIOZOL



Management System  
ISO 9001:2015

www.tuv.com  
ID 900019771



## Application Recommendation

**Application:** ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

**Dilution:** ELISA 1:10.000 - 1:50.000, Immunohistochemistry (IHC) 1:500 - 1:2,500, Western Blot (WB) 1:1.000 - 1:10.000

### Application Note:

Mouse IgG F(c) Antibody is suitable for immunoblotting (western or dot blot), ELISA, and immunohistochemistry requiring extremely low background levels, lot-to-lot consistency, high titer and specificity.

## Background Information

Anti-Mouse IgG F(c) generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of mouse IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity.

## Safety Information

This reagent contains Gentamicin Sulfate (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material. **DO NOT ADD SODIUM AZIDE!** Use of Sodium Azide as a preservative will substantially inhibit the enzyme activity of horseradish peroxidase!

## Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

### Manufacturer

**BIOZOL**  
Diagnostica Vertrieb GmbH  
Leipziger Straße 4  
85386 Eching

Phone +49 (89) 3799 666 6  
Fax +49 (89) 3799 666 99  
E-mail [info@biozol.de](mailto:info@biozol.de)  
[www.biozol.de](http://www.biozol.de)

[www.dianova.com](http://www.dianova.com)

Your Secondary Antibody Portal  
Powered by BIOZOL



Management System  
ISO 9001:2015



[www.tuv.com](http://www.tuv.com)  
ID 9000019771