

Product Datasheet / Instruction for use



SEC-182665 2023-02-23 1 / 2

Rabbit F(ab')2 anti-Dog IgG (F(ab')2)-HRPO, MinX none

General Information

Catalog Number: SEC-182665 Physical State: Lyophilized

Quantity: 20 mg Reconstitution Volume: 2.0 mL

Reconstitution Buffer: Restore with deionized water (or equivalent)

Shipping Conditions: at room temperature

Product Expiration Expiration date is one (1) year from date of

opening.

Antibody Host / Format

Host Species: Rabbit Clonality / Isotype / Clone: polyclonal Ig

Antibody Format: IgG F(ab')2 Fragment Concentration: 10.0 mg/mL by UV absorbance at 280

nm

Conjugation: Horseradish Peroxidase (HRP)

Maximum Excitation / Emission: /

Specificity

Target Species: Dog Immunogen: Dog IgG F(ab')2 fragment

Antibody Specificity: IgG (F(ab')2 This product is a F(ab')2 fragment of an IgG fraction antibody purified

from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Dog IgG, Dog IgG F(ab')2 and Dog Serum. No reaction was observed against Dog IgG F(c), anti-Rabbit IgG

F(c) or anti-Pepsin.

Minimal Cross Reactivity: none

Formulation, Transport and Storage

Storage Buffer: 0.01 M Sodium Phosphate,

Fragment)

0.15 M Sodium Chloride, pH 7.2

Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease

free

Preservatives: 0.01% (w/v) Gentamicin

Sulfate

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Storage Temperature: 4-8°C

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL
Diagnostica Vertrieb GmbH
Leipziger Straße 4
85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com

Your Secondary Antibody Portal Powered by BIOZOL



Management System ISO 9001:2015



www.tuv.com ID 9000019771



Product Datasheet / Instruction for use



SEC-182665 2023-02-23 2 / 2

Application Recommendation

Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections),

Immunohistochemistry (frozen sections), Western Blot

Dilution: ELISA 1:10.000 - 1:50.000, Immunohistochemistry (IHC) 1:500 - 1:2,500, Western Blot (WB) 1:1.000 - 1:10.000

Application Note:

This product has been assayed against 1.0 ug of Dog IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product.

Background Information

F(ab')2 Anti-Dog IgG F(ab')2 Peroxidase Antibody generated in rabbit detects Dog F(ab')2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Safety Information

This reagent contains Gentamicin Sulfate (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.DO NOT ADD SODIUM AZIDE! Use of Sodium Azide as a preservative will substantially inhibit the enzyme activity of horseradish peroxidase!

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.





