

Product Datasheet / Instruction for use



Goat F(ab')2 anti-Horse IgG (H+L)-FITC, MinX none

General Information

Catalog Number:	SEC-182681	Physical State:	Lyophilized
Quantity:	20 mg	Reconstitution Volume:	2.0 mL
		Reconstitution Buffer:	Restore with deionized water (or equivalent)
		Shipping Conditions:	at room temperature
		Product Expiration	Expiration date is one (1) year from date of opening.
Antibody Host /	Format		
Host Species:	Goat	Clonality / Isotype / Clon	e: polyclonal Ig
Antibody Format:	IgG F(ab')2 Fragment	Concentration:	
		Conjugation:	FITC (Fluoresceinisothiocyanat)
		Maximum Excitation / Emission: 492 nm / 520 nm	
Specificity			
Target Species:	Horse	Immunogen:	Horse IgG whole molecule
Antibody Specificity: IgG (H+L)		This product is a F(ab')2 fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and peps digestion followed by extensive dialysis against the buffer stated above Assay by immunoelectrophoresis resulted in a single precipitin arc again anti-fluorescein, anti-Goat Serum, Horse IgG and Horse Serum. No reaction was observed against anti-Goat IgG F(c) or anti-Pepsin.	
		digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Goa	ensive dialysis against the buffer stated above. phoresis resulted in a single precipitin arc again t Serum, Horse IgG and Horse Serum. No
		digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Goa	ensive dialysis against the buffer stated above. phoresis resulted in a single precipitin arc again at Serum, Horse IgG and Horse Serum. No painst anti-Goat IgG F(c) or anti-Pepsin.
Formulation, Tra	ansport and Storage	digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Goa reaction was observed ag	ensive dialysis against the buffer stated above. phoresis resulted in a single precipitin arc again at Serum, Horse IgG and Horse Serum. No painst anti-Goat IgG F(c) or anti-Pepsin.
Storage Buffer: 0 Phosphate, 0.15 M	·	digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Goa reaction was observed ag Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -2 thawing. Centrifuge proc room temperature. This	ensive dialysis against the buffer stated above. ohoresis resulted in a single precipitin arc again at Serum, Horse IgG and Horse Serum. No gainst anti-Goat IgG F(c) or anti-Pepsin. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and luct if not completely clear after standing at product is stable for several weeks at 4° C as ar
Storage Buffer: 0 Phosphate, 0.15 M Stabilizer: N).02 M Potassium 1 Sodium Chloride, pH 7.2	digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Goa reaction was observed ag Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -2 thawing. Centrifuge proc room temperature. This	ensive dialysis against the buffer stated above. ohoresis resulted in a single precipitin arc again at Serum, Horse IgG and Horse Serum. No gainst anti-Goat IgG F(c) or anti-Pepsin. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and luct if not completely clear after standing at

Dilution: FLISA 1:10.000 - 1:50.000, Flow cytometry 1:500 - 1:2,500, Immunofluorescence 1:1,000 - 1:5,000

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT

Management System ISO 9001:2015



Product Datasheet / Instruction for use



Application Note:

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.

Background Information

F(ab')2 Anti-Horse IgG Fluorescein Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Safety Information

The product contains no hazardous constituents. Standard Laboratory Practices should be followed when handling this material.

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer BIOZOL

Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL



Management System ISO 9001:2015