



# Rabbit F(ab')2 anti-Mouse IgG (F(ab')2)-HRPO, MinX none

# **General Information**

Catalog Number: SEC	C-182701	Physical State:	Lyophilized		
Quantity:	20 mg	<b>Reconstitution Volume:</b>	2.0 mL		
		<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)		
		Shipping Conditions:	at room temperature		
		Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / Form	nat				
Host Species: Ra	bbit	Clonality / Isotype / Clon	ne: polyclonal Ig		
Antibody Format: IgG	6 F(ab')2 Fragment	Concentration:	10.0 mg/mL by UV absorbance at 280 nm		
		Conjugation:	Horseradish Peroxidase (HRP)		
		Maximum Excitation / Emission: /			
Specificity					
Target Species:	Mouse	Immunogen:	Mouse IgG F(ab')2 fragment		
Antibody Specificity:	lgG (F(ab')2 Fragment)	This product is a F(ab')2 fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Mouse IgG, Mouse IgG F(ab')2 and Mouse Serum. No reaction was observed against Mouse IgG F(c), anti- Rabbit IgG F(c) or anti-Pepsin.			
		Minimal Cross Reactivity: none			
Formulation, Transpo	ort and Storage	l			
Storage Buffer: 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2		Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at			
Stabilizer:10 mg/mL Bovine SerumAlbumin (BSA) - Immunoglobulin and Proteasefree		room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. Storage Temperature: 4-8°C			
		-			

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.						
Manufacturer				Management System		
<b>BIOZOL</b> Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015		
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771		



#### Product Datasheet / Instruction for use



### **Application Recommendation**

**Application:** ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

Dilution: ELISA 1:10.000 - 1:50.000, Immunohistochemistry (IHC) 1:500 - 1:2,500, Western Blot (WB) 1:1.000 - 1:10.000

#### **Application Note:**

F(ab')2 Anti-Mouse IgG F(ab')2 Peroxidase Antibody has been assayed against 1.0 ug of Mouse IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:50,000 of the reconstitution concentration is suggested for this product.

# **Background Information**

F(ab')2 Anti-Mouse IgG F(ab')2 Peroxidase Antibody generated in rabbit detects Mouse F(ab')2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

# **Safety Information**

This reagent contains Gentamicin Sulfate (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.DO NOT ADD SODIUM AZIDE! Use of Sodium Azide as a preservative will substantially inhibit the enzyme activity of horseradish peroxidase!

## Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

# Manufacturer

BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT

Management System ISO 9001:2015