

Product Datasheet / Instruction for use



Rabbit F(ab')2 anti-Rat IgG (F(ab')2)-FITC, MinX none

General Information

Catalog Number: SEC-182710	Physical State:	Lyophilized 2.0 mL		
Quantity: 20 mg	Reconstitution Volume:			
	Reconstitution Buffer:	Restore with deionized water (or equivalent)		
	Shipping Conditions:	at room temperature		
	Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / Format				
Host Species: Rabbit	Clonality / Isotype / Clor	one: polyclonal Ig		
Antibody Format: IgG F(ab')2 Fragment	Concentration:	10.0 mg/mL by UV absorbance at 280 nm		
	Conjugation:	FITC (Fluoresceinisothiocyanat)		
	Maximum Excitation / E	Maximum Excitation / Emission: 492 nm / 520 nm		
Specificity	I			
Target Species: Rat	Immunogen:	Rat IgG F(ab')2 fragment		
Antibody Specificity: IgG (F(ab')2	This product is a F(ab')2 fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and peps digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc again anti-fluorescein, anti-Rabbit Serum, Rat IgG, Rat IgG F(ab')2 and Rat Serum. No reaction was observed against Rat IgG F(c), anti-Rabbit IgG F(c) or anti-Pepsin.			
Fragment)	delipidation, salt fraction digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Rab Serum. No reaction was	ation, ion exchange chromatography and pepsi tensive dialysis against the buffer stated above. phoresis resulted in a single precipitin arc again obit Serum, Rat IgG, Rat IgG F(ab')2 and Rat		
Fragment)	delipidation, salt fraction digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Rab Serum. No reaction was	ation, ion exchange chromatography and pepsi tensive dialysis against the buffer stated above. phoresis resulted in a single precipitin arc again obit Serum, Rat IgG, Rat IgG F(ab')2 and Rat observed against Rat IgG F(c), anti-Rabbit IgG		
Formulation, Transport and Storage	delipidation, salt fraction digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Rab Serum. No reaction was F(c) or anti-Pepsin.	ation, ion exchange chromatography and pepsi ensive dialysis against the buffer stated above. phoresis resulted in a single precipitin arc agains obit Serum, Rat IgG, Rat IgG F(ab')2 and Rat observed against Rat IgG F(c), anti-Rabbit IgG		
	delipidation, salt fraction digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Rab Serum. No reaction was F(c) or anti-Pepsin. Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -2	ation, ion exchange chromatography and pepsi tensive dialysis against the buffer stated above. phoresis resulted in a single precipitin arc again obit Serum, Rat IgG, Rat IgG F(ab')2 and Rat observed against Rat IgG F(c), anti-Rabbit IgG renone		
Formulation, Transport and Storage Storage Buffer: 0.01 M Sodium Phosphate,	 delipidation, salt fraction digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Rab Serum. No reaction was F(c) or anti-Pepsin. Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -2 thawing. Centrifuge proor room temperature. This 	ation, ion exchange chromatography and pepsi tensive dialysis against the buffer stated above. phoresis resulted in a single precipitin arc again obit Serum, Rat IgG, Rat IgG F(ab')2 and Rat observed against Rat IgG F(c), anti-Rabbit IgG r: none		
Formulation, Transport and Storage Storage Buffer: 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease	 delipidation, salt fraction digestion followed by ext Assay by immunoelectrop anti-fluorescein, anti-Rab Serum. No reaction was F(c) or anti-Pepsin. Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -2 thawing. Centrifuge proor room temperature. This 	ation, ion exchange chromatography and pepsi tensive dialysis against the buffer stated above. phoresis resulted in a single precipitin arc again obit Serum, Rat IgG, Rat IgG F(ab')2 and Rat observed against Rat IgG F(c), anti-Rabbit IgG restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and duct if not completely clear after standing at product is stable for several weeks at 4° C as ar		

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.				
Manufacturer				Management System
BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771



Product Datasheet / Instruction for use



Application: FLISA, Flow Cytometry, Immunofluorescence

Dilution: FLISA 1:10.000 - 1:50.000, Flow cytometry 1:500 - 1:2,500, Immunofluorescence 1:1,000 - 1:5,000

Application Note:

F(ab')2 Anti-Rat IgG F(ab')2 Fluorescein Antibody is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.

Background Information

F(ab')2 Anti-Rat IgG F(ab')2 Fluorescein Antibody generated in rabbit detects Rat F(ab')2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Safety Information

This reagent contains Thimerosal (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT

Management System ISO 9001:2015