

Product Datasheet / Instruction for use



# Goat IgG anti-Cat IgG (H+L)-Biotin, MinX none

## **General Information**

| Catalog Number: SEC-182755  | Physical State:   | Lyophilized  |  |  |  |
|---|---|--|--|--|--|
| Quantity: 2 mg  | Reconstitution Volume:  | 1.0 mL   |  |  |  |
|   | <b>Reconstitution Buffer:</b>   | Restore with deionized water (or equivalent)   |  |  |  |
|   | Shipping Conditions:  | at room temperature  |  |  |  |
|   | Product Expiration  | Expiration date is one (1) year from date of opening.  |  |  |  |
| Antibody Host / Format  |   |  |  |  |  |
| Host Species: Goat  | Clonality / Isotype / Clon  | ty / Isotype / Clone: polyclonal Ig  |  |  |  |
| Antibody Format: IgG whole molecule   | Concentration:  | 2.0 mg/mL by UV absorbance at 280 nn   |  |  |  |
|   | Conjugation:  | Biotin   |  |  |  |
|   | Maximum Excitation / Emission: /  |  |  |  |  |
| Specificity   |   |  |  |  |  |
| Target Species: Cat   | Immunogen:  | Cat IgG whole molecule   |  |  |  |
| Antibody Specificity: IgG (H+L)   | This product was prepared from monospecific antiserum by<br>immunoaffinity chromatography using Cat IgG coupled to agarose beads<br>followed by solid phase adsorption(s) to remove any unwanted<br>reactivities. Assay by immunoelectrophoresis resulted in a single<br>precipitin arc against anti-biotin, anti-Goat Serum, Cat IgG and Cat Serum |  |  |  |  |
|   | Minimal Cross Reactivity  | : none   |  |  |  |
| Formulation, Transport and Storage  | I   |  |  |  |  |
| Storage Buffer: 0.02 M Potassium<br>Phosphate, 0.15 M Sodium Chloride, pH 7.2 | contents and freeze at -2   | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and     |  |  |  |
| Stabilizer: None  |   | thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an |  |  |  |
| Preservatives: None   | undiluted liquid. Dilute o  | nly prior to immediate use.  |  |  |  |
|   | Storage Temperature:  | 4-8°C  |  |  |  |
| Application Recommendation  | I   |  |  |  |  |

Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

| This product is for In Vitro exp                          | erimental use only. Not for Ther   | apeutic or Diagnostic use.                        |              |                              |     |
|---|--|---|--------------|------------------------------|-----|
| Manufacturer  |  |   |              | Management<br>System         | 250 |
| BIOZOL<br>Diagnostica Vertrieb GmbH<br>Leipziger Straße 4 | Phone +49 (89) 3799 666 6<br>Fax +49 (89) 3799 666 99<br>E-mail info@biozol.de | www.dianova.com<br>Your Secondary Antibody Portal | TÜVRheinland | ISO 9001:2015                |     |
| 85386 Eching  | www.biozol.de  | Powered by BIOZOL                                 | ZERTIFIZIERT | www.tuv.com<br>ID 9000019771 |     |



#### Product Datasheet / Instruction for use



Dilution: ELISA 1:20.000 - 1:100.000, Immunohistochemistry (IHC) 1:1,000 - 1:5,000, Western Blot (WB) 1:2.000 -1:10.000

#### **Application Note:**

Secondary antibody reagents are ideal for ELISA, western blotting, Immunohistochemistry, Fluorescence Microscopy, Flow Cytometry as well as other antibody detection methods.

#### **Background Information**

Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. Anti-Cat IgG (H&L) antibody is ideal for investigators in Immunology, Cancer, and Microbiology research.

### Safety Information

The product contains no hazardous constituents. Standard Laboratory Practices should be followed when handling this material.

#### Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL **Diagnostica Vertrieb GmbH** Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL



Management System ISO 9001:2015