



Rabbit IgG anti-Hamster generally IgG (H+L)-Alk. Phos., MinX none

General Information

Quantity: 1 mg Reconstitution Volume: Reconstitution Buffer: Shipping Conditions: Wether Product Expiration Expiration Expiration Antibody Host / Format Clonality / Isotype / Clone: Ope Antibody Format: IgG whole molecule Concentration: Conjugation: Maximum Excitation / Emission Maximum Excitation / Emission Specificity Target Species: Hamster generally Immunogen: Gold Antibody Specificity: IgG (H+L) This product was prepared from immunoaffinity chromatograp coupled to agarose beads. Assistingle precipitin arc against an anti-Rabbit Serum, Golden Syrt Hamster Serum. Minimal Cross Reactivity: non serum. Formulation, Transport and Storage Storage Buffer: 0.05 M Tris Chloride, 0.15M Store vial at 4° C before openin at 4° C as an undiluted liquid.				
Reconstitution Buffer: Shipping Conditions: Wet Product Expiration Expiration ope Antibody Host / Format Host Species: Rabbit Antibody Format: IgG whole molecule Concentration: Concentration: Conjugation: Maximum Excitation / Emission Specificity Immunogen: Gold Target Species: Hamster generally Ins product was prepared from immunoaffinity chromatograp coupled to agarose beads. Assigned precipitin arc against an anti-Rabbit Serum, Golden Syrt Hamster Serum. Minimal Cross Reactivity: non Formulation, Transport and Storage Storage Buffer: 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.001M Zinc Chloride, 50% (v/v) Preservatives: 0.01% (w/v) Store vial at 4° C before openin at 4° C as an undiluted liquid.	d (sterile filtered)			
Shipping Conditions: Wett Product Expiration Expiration Antibody Host / Format Clonality / Isotype / Clone: Antibody Format: IgG whole molecule Antibody Format: IgG whole molecule Concentration: Conjugation: Maximum Excitation / Emission Specificity Target Species: Hamster generally Antibody Specificity: IgG (H+L) This product was prepared from immunoaffinity chromatograp coupled to agarose beads. Assistingle precipitin arc against an anti-Rabbit Serum, Golden Synthamster Serum. Minimal Cross Reactivity: Non Formulation, Transport and Storage Preservatives: 0.01% (w/v) 1 Sodium Chloride, 0.001M Magnesium Chloride, 50% (v/v) Store vial at 4° C before openinat 4° C as an undiluted liquid.				
Antibody Host / Format Product Expiration Expiration Expiration Host Species: Rabbit Clonality / Isotype / Clone: Antibody Format: IgG whole molecule Concentration: Specificity Conjugation: Maximum Excitation / Emission Specificity Target Species: Hamster generally Immunogen: Gold Antibody Specificity: IgG (H+L) This product was prepared from immunoaffinity chromatograp coupled to agarose beads. Assistingle precipitin arc against an anti-Rabbit Serum, Golden Syrt Hamster Serum. Minimal Cross Reactivity: non Formulation, Transport and Storage Storage Buffer: 0.05 M Tris Chloride, 0.15M Store vial at 4° C before openin at 4° C before openin at 4° C as an undiluted liquid.				
Antibody Host / Format Host Species: Rabbit Antibody Format: IgG whole molecule Concentration: Conjugation: Maximum Excitation / Emission Specificity Target Species: Hamster generally Antibody Specificity: IgG (H+L) This product was prepared from immunoaffinity chromatograp coupled to agarose beads. Assistingle precipitin arc against an anti-Rabbit Serum, Golden Syrt Hamster Serum. Minimal Cross Reactivity: non Formulation, Transport and Storage Storage Buffer: 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.001M Zinc Chloride, 50% (v/v) Givceroi: pH 8.0	ce			
Host Species: Rabbit Clonality / Isotype / Clone: Antibody Format: IgG whole molecule Concentration: Antibody Format: IgG whole molecule Conjugation: Maximum Excitation / Emission Maximum Excitation / Emission Specificity Target Species: Hamster generally Antibody Specificity: IgG (H+L) Immunogen: Gold Antibody Specificity: IgG (H+L) This product was prepared fro immunoaffinity chromatograp coupled to agarose beads. Assessingle precipitin arc against an anti-Rabbit Serum, Golden Syr Hamster Serum. Minimal Cross Reactivity: non Formulation, Transport and Storage Storage Buffer: 0.05 M Tris Chloride, 0.15M Preservatives: 0.01% (w/v) Store vial at 4° C before openin at 4° C as an undiluted liquid.	ration date is one (1) year from date of ning.			
Antibody Format: IgG whole molecule Concentration: Conjugation: Maximum Excitation / Emission Specificity Maximum Excitation / Emission Target Species: Hamster generally Immunogen: Gold Antibody Specificity: IgG (H+L) This product was prepared from immunoaffinity chromatograp coupled to agarose beads. Assessingle precipitin arc against an anti-Rabbit Serum, Golden Syrt Hamster Serum. Minimal Cross Reactivity: none Formulation, Transport and Storage Preservatives: 0.01% (w/v) Store vial at 4° C before openin at 4° C as an undiluted liquid.				
Conjugation: Maximum Excitation / Emission Specificity Target Species: Hamster generally Antibody Specificity: IgG (H+L) This product was prepared from immunoaffinity chromatograp coupled to agarose beads. Assigned precipitin arc against an anti-Rabbit Serum, Golden Syrt Hamster Serum. Minimal Cross Reactivity: non Formulation, Transport and Storage Storage Buffer: 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol: pH 8.0	polyclonal Ig			
Specificity Target Species: Hamster generally Antibody Specificity: IgG (H+L) This product was prepared fro immunoaffinity chromatograp coupled to agarose beads. Ass single precipitin arc against an anti-Rabbit Serum, Golden Syr Hamster Serum. Minimal Cross Reactivity: non Formulation, Transport and Storage Storage Buffer: 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol: pH 8.0	1.0 mg/mL by UV absorbance at 280 nr			
Specificity Target Species: Hamster generally Antibody Specificity: IgG (H+L) This product was prepared from immunoaffinity chromatograp coupled to agarose beads. Assigning precipitin arc against an anti-Rabbit Serum, Golden Syrt Hamster Serum. Minimal Cross Reactivity: non Formulation, Transport and Storage Storage Buffer: 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol: pH 8.0	Alkaline Phosphatase			
Target Species:Hamster generallyImmunogen:GoldAntibody Specificity:IgG (H+L)This product was prepared fro immunoaffinity chromatograp coupled to agarose beads. Ass single precipitin arc against an 	Maximum Excitation / Emission: /			
Antibody Specificity: IgG (H+L) This product was prepared from immunoaffinity chromatograp coupled to agarose beads. Assigned precipitin arc against an anti-Rabbit Serum, Golden Syrt Hamster Serum. Minimal Cross Reactivity: None Formulation, Transport and Storage Preservatives: 0.01% (w/v) Store vial at 4° C before opening at 4° C as an undiluted liquid.				
immunoaffinity chromatograp coupled to agarose beads. Ass single precipitin arc against an anti-Rabbit Serum, Golden Syr Hamster Serum. Minimal Cross Reactivity: non Formulation, Transport and Storage Storage Buffer: 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol: pH 8.0	en Syrian Hamster IgG, whole molecule			
Formulation, Transport and StorageStorage Buffer:0.05 M Tris Chloride, 0.15MSodium Chloride, 0.001M MagnesiumPreservatives:0.01% (w/v)Chloride, 0.0001M Zinc Chloride, 50% (v/v)Store vial at 4° C before openin at 4° C as an undiluted liquid.	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Golden Syrian Hamster IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Golden Syrian Hamster IgG, and Golden Syrian Hamster Serum.			
Storage Buffer:0.05 M Tris Chloride, 0.15MPreservatives:0.01% (w/v)Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol: pH 8.0Store vial at 4° C before openin at 4° C as an undiluted liquid.	Minimal Cross Reactivity: none			
Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol: pH 8.0 Store vial at 4° C before openin at 4° C as an undiluted liquid.				
Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol: pH 8.0 Store vial at 4° C before openin at 4° C as an undiluted liquid.	Preservatives: 0.01% (w/v) Sodium Azide			
	Store vial at 4° C before opening. DO NOT FREEZE. This product is stab at 4° C as an undiluted liquid. Dilute only prior to immediate use. Freezing alkaline phosphatase conjugates will result in a substantial loss			
Stabilizer:10 mg/mL Bovine Serumof enzymatic activity.Albumin (BSA) - Immunoglobulin and Protease freeStorage Temperature:4-8°				
Application Recommendation				

Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.					
Manufacturer				Management System	
BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015	
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771	



Product Datasheet / Instruction for use



Dilution: ELISA 1:5.500, Immunohistochemistry (IHC) 1:200 - 1:1,000, Western Blot (WB) 1:200 - 1:1.000

Application Note:

Anti-Golden Syrian Hamster IgG Alk Phos conjugate is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring lot-to-lot consistency.

Background Information

Anti-Golden Syrian Hamster IgG Alkaline Phosphatase Antibody generated in rabbit detects Golden Syrian Hamster IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Safety Information

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL



Management System ISO 9001:2015

www.tuv.com ID 9000019771