

Product Datasheet / Instruction for use



# Goat IgG anti-Human IgG (Fc)-HRPO, MinX Ms

## **General Information**

Catalog Number: SEC-182993	Physical State:	Lyophilized		
Quantity: 1 mg	Reconstitution Volume:	1.0 mL		
	<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)		
	Shipping Conditions:	at room temperature		
	Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / Format				
Host Species: Goat	Clonality / Isotype / Clor	ne: polyclonal Ig		
Antibody Format: IgG whole molecule	Concentration:	1.0 mg/mL by UV absorbance at 280 nr		
	Conjugation:	Horseradish Peroxidase (HRP)		
	Maximum Excitation / Emission: /			
Specificity	I			
Target Species: Human	Immunogen:	Human IgG F(c) fragment		
Antibody Specificity: IgG (Fc Fragment)	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to ag beads followed by solid phase adsorption(s) to remove any unw reactivities. Assay by immunoelectrophoresis resulted in a sing precipitin arc against anti-Peroxidase, anti-Goat Serum, Human Human IgG F(c) and Human Serum. No reaction was observed a Human IgG F(ab)or Mouse serum proteins.			
	Minimal Cross Reactivity	Minimal Cross Reactivity: Mouse		
Formulation, Transport and Storage	I			
Storage Buffer:0.02 M PotassiumPhosphate,0.15 M Sodium Chloride,Phosphate,0.15 M Sodium Chloride,Phosphate,10 mg/mL Bovine SerumAlbumin (BSA) - Immunoglobulin and Proteasefree	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.			
<b>Preservatives:</b> 0.01% (w/v) Gentamicin Sulfate	Storage Temperature: 4-8°C			
Application Recommendation	I			

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.							
Manufacturer				Management System			
BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com	TÜVRheinland	ISO 9001:2015			
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771			



#### Product Datasheet / Instruction for use



Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

**Dilution:** ELISA 1:200.000, Immunohistochemistry (IHC) 1:500 - 1:2,500, Western Blot (WB) 1:1.000 - 1:10.000

#### **Application Note:**

Anti-Human IgG F(c) Peroxidase conjugate is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring lot-to-lot consistency.

#### **Background Information**

Anti-Human IgG F(c) Peroxidase Conjugated generated in goat detects Human F(c). A proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of human IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(c) Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

### Safety Information

This reagent contains Gentamicin Sulfate (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.DO NOT ADD SODIUM AZIDE! Use of Sodium Azide as a preservative will substantially inhibit the enzyme activity of horseradish peroxidase!

### Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL **Diagnostica Vertrieb GmbH** Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT ID 9000019771

Management ISO 9001:2015

System