



Goat IgG anti-Human IgG (H+L)-HRPO, MinX Bo,Ck,Go,Gp,Hm,Ho,Ms,Rb,Rt,Sh

General Information

Catalog Number:	SEC-182994	Physical State:	Lyophilized		
Quantity:	1 mg	Reconstitution Volume:	1.0 mL		
		Reconstitution Buffer:	Restore with deionized water (or equivalent)		
		Shipping Conditions:	at room temperature		
		Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / F	Format				
Host Species:	Goat	Clonality / Isotype / Clor	ne: polyclonal Ig		
Antibody Format:	IgG whole molecule	Concentration:	1.0 mg/mL by UV absorbance at 280 nm		
		Conjugation:	Horseradish Peroxidase (HRP)		
		Maximum Excitation / Emission: /			
Specificity					
Target Species:	Human	Immunogen:	Human IgG whole molecule		
Antibody Specificity: IgG (H+L)		HRP Conjugated Secondary Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Human IgG and Human Serum. No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Mouse, Rabbit, Rat and Sheep Serum Proteins.			
		Minimal Cross Reactivity: Bovine, Chicken, Goat, Guinea Pig, Hamster generally, Horse, Mouse, Rabbit, Rat, Sheep			
Formulation, Trai	nsport and Storage				
Storage Buffer:0.02 M PotassiumPhosphate,0.15 M Sodium Chloride,Phosphate,0.15 M Sodium Chloride,Stabilizer:10 mg/mL Bovine SerumAlbumin (BSA) - Immunoglobulin and Proteasefree		Store Secondary Antibody at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge HRP Conjugated Secondary Antibody if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.			
Preservatives: 0. Sulfate	01% (w/v) Gentamicin	Storage Temperature: 4-8°C			

This product is for In Vitro exp	perimental use only. Not for Ther	apeutic or Diagnostic use.			
Manufacturer				Management System	
BIOZOL Diagnostica Vertrieb GmbH	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015	
Leipziger Straße 4 85386 Eching	E-mail info@biozol.de www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771	



Product Datasheet / Instruction for use



5201

Application Recommendation

Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

Dilution: ELISA 1:200.000, Immunohistochemistry (IHC) 1:500 - 1:2,500, Western Blot (WB) 1:1.000 - 1:5.000

Application Note:

Peroxidase Secondary antibody reagents are ideal for western blotting, Immunohistochemistry and ELISA as well as other antibody detection methods

Background Information

Anti-Human IgG (H&L) Peroxidase generated in goat detects human Immunoglobulin G (IgG), both heavy and light chains of the antibody molecule are present. It is a protein complex composed of four peptide chains — two identical heavy chains and two identical light chains arranged in a Y-shape typical of antibody monomers. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins in humans, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Safety Information

This reagent contains Gentamicin Sulfate (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.DO NOT ADD SODIUM AZIDE! Use of Sodium Azide as a preservative will substantially inhibit the enzyme activity of horseradish peroxidase!

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT

Management System ISO 9001:2015