

Product Datasheet / Instruction for use



# Goat IgG anti-Mouse IgG2a (Fc)-ATTO 655, MinX none

## **General Information**

Catalog Number: SE	C-183221	Physical State: Lyophilized			
Quantity:	0,5 mg	Reconstitution Volume:	500 μL		
		<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent) at room temperature		
		Shipping Conditions:			
		Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / Forn	nat				
Host Species: Go	pat	Clonality / Isotype / Clone: polyclonal Ig			
Antibody Format: IgC	G whole molecule	Concentration:	1.0 mg/mL by UV absorbance at 280 nm		
		Conjugation:	ATTO 655		
		Maximum Excitation / Emission: 662 nm / 680 nm			
Specificity					
Target Species:	Mouse	Immunogen:	Mouse IgG2a heavy chain		
Antibody Specificity:	lgG2a (lgG Fc	MOUSE IgG2a Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse Serum and Mouse IgG2a. Specificity was confirmed by ELISA at less than 1% cross-reactivity against other mouse heavy or light chain isotypes. No reaction was observed against Bovine, Human, and Rabbit Serum Proteins. Specificity was confirmed by ELISA at less than 1% of target signal.			
Anibody Specificity.	gamma 2a)	immunoaffinity chromato followed by solid phase a reactivities. Assay by im- precipitin arc against anti Specificity was confirmed other mouse heavy or ligl against Bovine, Human, a	ography using antigens coupled to agarose beac dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a. I by ELISA at less than 1% cross-reactivity agains ht chain isotypes. No reaction was observed and Rabbit Serum Proteins. Specificity was		
	gamma 2a)	immunoaffinity chromato followed by solid phase a reactivities. Assay by im- precipitin arc against anti Specificity was confirmed other mouse heavy or ligl against Bovine, Human, a	ography using antigens coupled to agarose bead dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a. I by ELISA at less than 1% cross-reactivity agains ht chain isotypes. No reaction was observed nd Rabbit Serum Proteins. Specificity was s than 1% of target signal.		
Formulation, Transpo	gamma 2a)	immunoaffinity chromato followed by solid phase a reactivities. Assay by imp precipitin arc against anti Specificity was confirmed other mouse heavy or ligh against Bovine, Human, a confirmed by ELISA at les	ography using antigens coupled to agarose bead dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a. I by ELISA at less than 1% cross-reactivity agains ht chain isotypes. No reaction was observed nd Rabbit Serum Proteins. Specificity was s than 1% of target signal.		
	gamma 2a) ort and Storage M Potassium	immunoaffinity chromato followed by solid phase a reactivities. Assay by im- precipitin arc against anti Specificity was confirmed other mouse heavy or ligh against Bovine, Human, a confirmed by ELISA at les <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -20	ography using antigens coupled to agarose beac dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a. by ELISA at less than 1% cross-reactivity agains ht chain isotypes. No reaction was observed ind Rabbit Serum Proteins. Specificity was s than 1% of target signal. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and		
Formulation, Transpo Storage Buffer: 0.02 N Phosphate, 0.15 M Sodi	gamma 2a) ort and Storage M Potassium ium Chloride, pH 7.2 g/mL Bovine Serum	immunoaffinity chromato followed by solid phase a reactivities. Assay by im- precipitin arc against anti Specificity was confirmed other mouse heavy or ligi against Bovine, Human, a confirmed by ELISA at les <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge proc room temperature. This	bgraphy using antigens coupled to agarose beac dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single -Goat Serum, Mouse Serum and Mouse IgG2a. I by ELISA at less than 1% cross-reactivity agains ht chain isotypes. No reaction was observed ind Rabbit Serum Proteins. Specificity was is than 1% of target signal. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and fuct if not completely clear after standing at		
Formulation, Transports Storage Buffer: 0.02 M Phosphate, 0.15 M Sodi Stabilizer: 10 mg Albumin (BSA) - Immun- free	gamma 2a) ort and Storage M Potassium ium Chloride, pH 7.2 g/mL Bovine Serum	immunoaffinity chromato followed by solid phase a reactivities. Assay by im- precipitin arc against anti Specificity was confirmed other mouse heavy or ligi against Bovine, Human, a confirmed by ELISA at les <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge proc room temperature. This	<ul> <li>bgraphy using antigens coupled to agarose beact dsorption(s) to remove any unwanted munoelectrophoresis resulted in a single</li> <li>Goat Serum, Mouse Serum and Mouse IgG2a.</li> <li>by ELISA at less than 1% cross-reactivity agains the chain isotypes. No reaction was observed and Rabbit Serum Proteins. Specificity was s than 1% of target signal.</li> <li>: none</li> <li>restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and luct if not completely clear after standing at product is stable for several weeks at 4° C as an analysis.</li> </ul>		

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.						
Manufacturer				Management System		
BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015		
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771		



#### Product Datasheet / Instruction for use



Application: FLISA, Flow Cytometry, Immunofluorescence, Western Blot

Dilution: FLISA >1:20.000, Flow cytometry 1:500 - 1:2,500, Immunofluorescence >1:5,000, Western Blot (WB) 1:4.000 - 1:20.000

#### **Application Note:**

Mouse IgG2a secondary antibody is available in a variety of formats. ATTO 655 Conjugated antibodies are designed for STED microscopy, FRET, immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Anti-Mouse IgG2a secondary antibody is suitable for ELISA, Immunohistochemistry western blotting as well as other anti IgG2a antibody based assays. The emission spectra for this ATTO conjugate matches the principle output wavelengths of most common fluorescence instrumentation.

#### **Background Information**

Anti-Mouse IgG2a ATTO 655 Antibody generated in goat detects reactivity to Mouse IgG2a (Gamma 2a chain). Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. IgG2, the second largest of IgG isotypes, comprises almost 25% of IgG and has a low affinity for binding to the Fc receptor of phagocytic cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

### **Safety Information**

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

#### Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT Www.tuv.com ID 900019771

Management System ISO 9001:2015