

Product Datasheet / Instruction for use



SEC-183227 2023-02-23 1 / 2

Goat IgG anti-Mouse IgG (Fc)-Biotin, MinX none

General Information

Catalog Number: SEC-183227 Physical State: Lyophilized

Quantity: 2 mg Reconstitution Volume: 1.0 mL

Reconstitution Buffer: Restore with deionized water (or equivalent)

Shipping Conditions: at room temperature

Product Expiration Expiration date is one (1) year from date of

opening.

Antibody Host / Format

Host Species: Goat Clonality / Isotype / Clone: polyclonal Ig

Antibody Format: IgG whole molecule Concentration: 2.0 mg/mL by UV absorbance at 280 nm

Conjugation: Biotin

Maximum Excitation / Emission: /

Specificity

Antibody Specificity:

Target Species: Mouse Immunogen: Anti-mouse IgG F(c) was produced by repeated

immunization with mouse IgG F(c) fragment in

goat.

Anti-Mouse IgG F(c) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG F(ab).

Minimal Cross Reactivity: none

Formulation, Transport and Storage

Storage Buffer: 0.01 M Sodium Phosphate,

0.15 M Sodium Chloride, pH 7.2

Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease

free

Preservatives: 0.01% (w/v) Thimerosal

Application Recommendation

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Storage Temperature: 4-8°C

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

IgG (Fc Fragment)

Manufacturer

BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de



Your Secondary Antibody Portal Powered by BIOZOL



Management System ISO 9001:2015



www.tuv.com ID 9000019771





Product Datasheet / Instruction for use

SEC-183227 2023-02-23 2 / 2

Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections),

Immunohistochemistry (frozen sections), Western Blot

Dilution: ELISA 1:20.000 - 1:100.000, Immunohistochemistry (IHC) 1:1,000 - 1:5,000, Western Blot (WB) 1:2.000 -

1:10.000

Application Note:

Anti-Mouse IgG F(c) Antibody has been assayed against 1.0 ug of Mouse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:40,000 to 1:130,000 of the reconstitution concentration is suggested for Anti-Mouse IgG F(c) Antibody.

Background Information

Anti-Mouse IgG F(c) Antibody generated in goat detects specifically mouse IgG F(c). It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of mouse IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity. Anti-Mouse IgG F(c) Biotin conjugation antibody is ideal for investigators involved in Serum Protein Component research, Cancer, Immunology, and Microbiology research.

Safety Information

This reagent contains Thimerosal (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.







