

## Donkey IgG anti-Rabbit IgG (H+L)-Biotin, MinX none

### General Information

|                                   |  |
|-----------------------------------|--|
| <b>Catalog Number:</b> SEC-183434 | <b>Physical State:</b> Lyophilized   |
| <b>Quantity:</b> 2 mg             | <b>Reconstitution Volume:</b> 1.0 mL   |
|                                   | <b>Reconstitution Buffer:</b> Restore with deionized water (or equivalent)       |
|                                   | <b>Shipping Conditions:</b> at room temperature                                  |
|                                   | <b>Product Expiration:</b> Expiration date is one (1) year from date of opening. |

### Antibody Host / Format

|  |  |
|--|--|
| <b>Host Species:</b> Donkey                | <b>Clonality / Isotype / Clone:</b> polyclonal Ig          |
| <b>Antibody Format:</b> IgG whole molecule | <b>Concentration:</b> 2.0 mg/mL by UV absorbance at 280 nm |
|  | <b>Conjugation:</b> Biotin                                 |
|  | <b>Maximum Excitation / Emission:</b> /                    |

### Specificity

|  |   |
|--|---|
| <b>Target Species:</b> Rabbit          | <b>Immunogen:</b> Rabbit IgG whole molecule   |
| <b>Antibody Specificity:</b> IgG (H+L) | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Donkey Serum, Rabbit IgG and Rabbit Serum. |
|  | <b>Minimal Cross Reactivity:</b> none   |

### Formulation, Transport and Storage

|   |   |
|---|---|
| <b>Storage Buffer:</b> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2         | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| <b>Stabilizer:</b> 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |   |
| <b>Preservatives:</b> 0.01% (w/v) Sodium Azide  |   |
|   | <b>Storage Temperature:</b> 4-8°C   |

### Application Recommendation

**Application:** ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

#### Manufacturer

**BIOZOL**  
Diagnostica Vertrieb GmbH  
Leipziger Straße 4  
85386 Eching

Phone +49 (89) 3799 666 6  
Fax +49 (89) 3799 666 99  
E-mail info@biozol.de  
www.biozol.de

[www.dianova.com](http://www.dianova.com)

Your Secondary Antibody Portal  
Powered by BIOZOL



Management System  
ISO 9001:2015

www.tuv.com  
ID 900019771



**Dilution:** ELISA 1:300.000, Immunohistochemistry (IHC) 1:500 - 1:3000, Western Blot (WB) 1:3.000 - 1:15.000

**Application Note:**

Secondary antibody reagents are ideal for ELISA, western blotting, Immunohistochemistry, Fluorescence Microscopy, Flow Cytometry as well as other antibody detection methods.

**Background Information**

Anti-Rabbit IgG (H&L) Biotin Antibody generated in donkey detects reactivity to Rabbit IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both the Heavy and Light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

**Safety Information**

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

**Disclaimer**

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

**Manufacturer**

**BIOZOL**  
Diagnostica Vertrieb GmbH  
Leipziger Straße 4  
85386 Eching

Phone +49 (89) 3799 666 6  
Fax +49 (89) 3799 666 99  
E-mail info@biozol.de  
www.biozol.de

[www.dianova.com](http://www.dianova.com)

Your Secondary Antibody Portal  
Powered by BIOZOL



Management System  
ISO 9001:2015



www.tuv.com  
ID 900019771