

Product Datasheet / Instruction for use



Goat IgG anti-Ferret IgA-HRPO, MinX none

General Information

Catalog Number: SEC	C-183631	Physical State:	Lyophilized
Quantity:	1 mg	Reconstitution Volume:	1.0 mL
		Reconstitution Buffer:	Restore with deionized water (or equivalent)
		Shipping Conditions:	at room temperature
		Product Expiration	Expiration date is one (1) year from date of opening.
Antibody Host / Form	nat		
Host Species: Go	at	Clonality / Isotype / Clon	e: polyclonal Ig
Antibody Format:	6 whole molecule	Concentration:	1.0 mg/ml by UV absorbance at 280 nm
		Conjugation:	Horseradish Peroxidase (HRP)
		Maximum Excitation / Emission: /	
Specificity			
Target Species: Antibody Specificity:	Ferret IgA (Alpha-Kette /	Immunogen:	Anti-Ferret IgA was produced by repeated immunization with ferret IgA alpha heavy chai in goat.
	Fc-Fragment)	antiserum by immunoaffi agarose beads followed b unwanted reactivities. A single precipitin arc again and Ferret Serum. Speci	by solid phase adsorption(s) to remove any assay by immunoelectrophoresis resulted in a
		antiserum by immunoaffi agarose beads followed b unwanted reactivities. A single precipitin arc again and Ferret Serum. Speci	inity chromatography using Ferret IgA coupled to by solid phase adsorption(s) to remove any assay by immunoelectrophoresis resulted in a list anti-Peroxidase, anti-Goat Serum, Ferret IgA ficity was confirmed by ELISA at less than 1% ther Ferret heavy or light chain isotypes.
Formulation, Transpo		antiserum by immunoaffi agarose beads followed b unwanted reactivities. A single precipitin arc again and Ferret Serum. Speci cross reactivity against ot	inity chromatography using Ferret IgA coupled to by solid phase adsorption(s) to remove any assay by immunoelectrophoresis resulted in a list anti-Peroxidase, anti-Goat Serum, Ferret IgA ficity was confirmed by ELISA at less than 1% ther Ferret heavy or light chain isotypes.
Formulation, Transpo Storage Buffer: 0.02 M Phosphate, 0.15 M Sodi	ort and Storage 1 Potassium	antiserum by immunoaffi agarose beads followed b unwanted reactivities. A single precipitin arc again and Ferret Serum. Speci cross reactivity against ot Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -20	inity chromatography using Ferret IgA coupled to by solid phase adsorption(s) to remove any assay by immunoelectrophoresis resulted in a list anti-Peroxidase, anti-Goat Serum, Ferret IgA ficity was confirmed by ELISA at less than 1% ther Ferret heavy or light chain isotypes. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and
Storage Buffer: 0.02 N Phosphate, 0.15 M Sodi	ort and Storage A Potassium um Chloride, pH 7.2 /mL Bovine Serum	antiserum by immunoaffi agarose beads followed b unwanted reactivities. A single precipitin arc again and Ferret Serum. Speci- cross reactivity against ot Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge prod temperature. This produce	inity chromatography using Ferret IgA coupled in by solid phase adsorption(s) to remove any assay by immunoelectrophoresis resulted in a list anti-Peroxidase, anti-Goat Serum, Ferret IgA ficity was confirmed by ELISA at less than 1% ther Ferret heavy or light chain isotypes. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and
Storage Buffer: 0.02 M Phosphate, 0.15 M Sodi Stabilizer: 10 mg, Albumin (BSA) - Immuno free	ort and Storage A Potassium um Chloride, pH 7.2 /mL Bovine Serum	antiserum by immunoaffi agarose beads followed b unwanted reactivities. A single precipitin arc again and Ferret Serum. Speci- cross reactivity against ot Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge prod temperature. This produce	inity chromatography using Ferret IgA coupled to by solid phase adsorption(s) to remove any assay by immunoelectrophoresis resulted in a list anti-Peroxidase, anti-Goat Serum, Ferret IgA ficity was confirmed by ELISA at less than 1% ther Ferret heavy or light chain isotypes. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and uct if not completely clear after standing at roo ct is stable for several weeks at 4° C as an

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use. Management System Manufacturer BIOZOL Phone +49 (89) 3799 666 6 ISO 9001:2015 7 www.dianova.com Diagnostica Vertrieb GmbH Fax +49 (89) 3799 666 99 TÜVRheinland Your Secondary Antibody Portal Leipziger Straße 4 E-mail info@biozol.de ZERTIFIZIERT www.tuv.com ID 9000019771 85386 Eching Powered by BIOZOL www.biozol.de



Product Datasheet / Instruction for use



Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

Dilution: ELISA 1:10.000 - 1:50.000, Immunohistochemistry (IHC) 1:500 - 1:2,500, Western Blot (WB) 1:1.000 - 1:5.000

Application Note:

Anti-Ferret IgA Peroxidase conjugate is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring lot-to-lot consistency. Ferret IgA (alpha chain) antibody has been assayed against 1.0 ug of Ferret IgA in a standard capture ELISA using ABTS (2,2'-azino-bis-[3ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000 of the reconstitution concentration is suggested for Anti-Ferret IgA (alpha chain) Antibody.

Background Information

Anti-Ferret IgA Peroxidase Antibody generated in goat detects immunoglobulin A from ferret. Immunoglobulin A (IgA) is an antibody that plays a critical role in mucosal immunity. IgA has two subclasses (IgA1 and IgA2) and can exist in a dimeric form called secretory IgA (sIgA). Anti-Ferret IgA (alpha chain) Antibody is ideal for investigators in Cancer, Immunology, and Microbiology research.

Safety Information

This reagent contains Thimerosal (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.DO NOT ADD SODIUM AZIDE! Use of Sodium Azide as a preservative will substantially inhibit the enzyme activity of horseradish peroxidase!

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL **Diagnostica Vertrieb GmbH** Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL



Management System ISO 9001:2015