

## Rabbit F(ab')<sub>2</sub> anti-Horse IgG (H+L)-Alk. Phos., MinX none

### General Information

|                                   |  |
|-----------------------------------|--|
| <b>Catalog Number:</b> SEC-183715 | <b>Physical State:</b> Liquid (sterile filtered)                                 |
| <b>Quantity:</b> 0,5 mg           | <b>Reconstitution Volume:</b>  |
|                                   | <b>Reconstitution Buffer:</b>  |
|                                   | <b>Shipping Conditions:</b> Wet Ice  |
|                                   | <b>Product Expiration:</b> Expiration date is one (1) year from date of opening. |

### Antibody Host / Format

|  |  |
|--|--|
| <b>Host Species:</b> Rabbit                              | <b>Clonality / Isotype / Clone:</b> polyclonal Ig          |
| <b>Antibody Format:</b> IgG F(ab') <sub>2</sub> Fragment | <b>Concentration:</b> 0.5 mg/mL by UV absorbance at 280 nm |
|  | <b>Conjugation:</b> Alkaline Phosphatase                   |
|  | <b>Maximum Excitation / Emission:</b> /                    |

### Specificity

|  |   |
|--|---|
| <b>Target Species:</b> Horse           | <b>Immunogen:</b> Horse IgG whole molecule  |
| <b>Antibody Specificity:</b> IgG (H+L) | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase, anti-Rabbit Serum, Horse IgG and Horse Serum. No reaction was observed against anti-Pepsin or anti-Rabbit IgG F(c). |
|  | <b>Minimal Cross Reactivity:</b> none   |

### Formulation, Transport and Storage

|  |  |
|--|--|
| <b>Storage Buffer:</b> 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol; pH 8.0 | Store vial at 4° C before opening. DO NOT FREEZE. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic activity. |
| <b>Stabilizer:</b> None  | <b>Storage Temperature:</b> 4-8°C  |

**Preservatives:** None

### Application Recommendation

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

#### Manufacturer

**BIOZOL**  
Diagnostica Vertrieb GmbH  
Leipziger Straße 4  
85386 Eching

Phone +49 (89) 3799 666 6  
Fax +49 (89) 3799 666 99  
E-mail info@biozol.de  
www.biozol.de

[www.dianova.com](http://www.dianova.com)

Your Secondary Antibody Portal  
Powered by BIOZOL



Management System  
ISO 9001:2015

www.tuv.com  
ID 9000019771



**Application:** ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

**Dilution:** ELISA 1:1.000-1:15.000, Immunohistochemistry (IHC) 1:250-1:500, Western Blot (WB) 1:500-1:1.000

**Application Note:**

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. This product has been assayed against 1.0 µg of Horse IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.

**Background Information**

F(ab')<sub>2</sub> Anti-Horse IgG Alkaline Phosphatase Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)<sub>2</sub> fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)<sub>2</sub> fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab)<sub>2</sub> fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')<sub>2</sub> Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

**Safety Information**

The product contains no hazardous constituents. Standard Laboratory Practices should be followed when handling this material.

**Disclaimer**

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

**Manufacturer**

**BIOZOL**  
Diagnostica Vertrieb GmbH  
Leipziger Straße 4  
85386 Eching

Phone +49 (89) 3799 666 6  
Fax +49 (89) 3799 666 99  
E-mail info@biozol.de  
www.biozol.de

[www.dianova.com](http://www.dianova.com)

Your Secondary Antibody Portal  
Powered by BIOZOL



Management System  
ISO 9001:2015



www.tuv.com  
ID 9000019771