

Product Datasheet / Instruction for use



Goat F(ab')2 anti-Human IgA-FITC, MinX none

General Information

Catalog Number: SE	C-183731	Physical State:	Lyophilized		
Quantity:	1 mg	Reconstitution Volume:	1.0 mL		
		Reconstitution Buffer:	Restore with deionized water (or equivalent) at room temperature		
		Shipping Conditions:			
		Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / Forn	nat				
Host Species: Go	pat	Clonality / Isotype / Clone: polyclonal Ig			
Antibody Format: Igo	G F(ab')2 Fragment	Concentration:	1.5 mg/mL by UV absorbance at 280 nr		
		Conjugation:	FITC (Fluoresceinisothiocyanat)		
		Maximum Excitation / Er	laximum Excitation / Emission:492 nm / 520 nm		
Specificity					
Target Species:	Human	Immunogen:	Human IgA alpha heavy chain		
Antibody Specificity:	IgA (Alpha-Kette /	This product was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose bea followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti- fluorescein, anti-Goat Serum, Human IgA and Human Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c). Specificity was confirmed by ELISA at less than 1% cross-reactivity against other human heavy or light chain isotypes.			
	Fc-Fragment)	followed by solid phase a reactivities, pepsin digest immunoelectrophoresis r fluorescein, anti-Goat Ser was observed against ant confirmed by ELISA at les	dsorption(s) to remove any unwanted tion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- rum, Human IgA and Human Serum. No reactio ti-Pepsin or anti-Goat IgG F(c). Specificity was s than 1% cross-reactivity against other human		
		followed by solid phase a reactivities, pepsin digest immunoelectrophoresis r fluorescein, anti-Goat Ser was observed against ant confirmed by ELISA at les	dsorption(s) to remove any unwanted tion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- rum, Human IgA and Human Serum. No reactio ti-Pepsin or anti-Goat IgG F(c). Specificity was s than 1% cross-reactivity against other human pes.		
Formulation, Transpo		followed by solid phase a reactivities, pepsin digest immunoelectrophoresis r fluorescein, anti-Goat Ser was observed against ant confirmed by ELISA at les heavy or light chain isoty	dsorption(s) to remove any unwanted tion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- rum, Human IgA and Human Serum. No reactio ti-Pepsin or anti-Goat IgG F(c). Specificity was s than 1% cross-reactivity against other human pes.		
Formulation, Transpo Storage Buffer: 0.01 M 0.25 M Sodium Chloride	ort and Storage И Sodium Phosphate,	followed by solid phase a reactivities, pepsin digest immunoelectrophoresis r fluorescein, anti-Goat Ser was observed against ant confirmed by ELISA at les heavy or light chain isoty Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -2	dsorption(s) to remove any unwanted tion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- rum, Human IgA and Human Serum. No reactio ti-Pepsin or anti-Goat IgG F(c). Specificity was s than 1% cross-reactivity against other human pes. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and		
Storage Buffer: 0.01 N 0.25 M Sodium Chloride	ort and Storage M Sodium Phosphate, e, pH 7.2 g/mL Bovine Serum	followed by solid phase a reactivities, pepsin digest immunoelectrophoresis r fluorescein, anti-Goat Ser was observed against ant confirmed by ELISA at les heavy or light chain isoty Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -2 thawing. Centrifuge proc room temperature. This undiluted liquid. Dilute o	dsorption(s) to remove any unwanted tion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- rum, Human IgA and Human Serum. No reaction ti-Pepsin or anti-Goat IgG F(c). Specificity was s than 1% cross-reactivity against other human pes. : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and duct if not completely clear after standing at product is stable for several weeks at 4° C as ar only prior to immediate use.		
Storage Buffer: 0.01 M 0.25 M Sodium Chloride Stabilizer: 15 mg Albumin (BSA) - Immun free	ort and Storage M Sodium Phosphate, e, pH 7.2 g/mL Bovine Serum	followed by solid phase a reactivities, pepsin digest immunoelectrophoresis r fluorescein, anti-Goat Ser was observed against ant confirmed by ELISA at les heavy or light chain isoty Minimal Cross Reactivity Store vial at 4° C prior to contents and freeze at -2 thawing. Centrifuge proc room temperature. This	dsorption(s) to remove any unwanted tion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- rum, Human IgA and Human Serum. No reaction ti-Pepsin or anti-Goat IgG F(c). Specificity was s than 1% cross-reactivity against other human pes. the none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and duct if not completely clear after standing at product is stable for several weeks at 4° C as ar		

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.						
Manufacturer				Management System		
BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015		
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771		



Product Datasheet / Instruction for use



Application: Flow Cytometry, Immunofluorescence

Dilution: Flow cytometry 1:500-1:2.500, Immunofluorescence 1:1.000-1:5.000

Application Note:

F(ab')2 Anti-Human IgA Fluorescein Antibody is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lotto-lot consistency, high titer and specificity.

Background Information

F(ab')2 Anti-Human IgA Fluorescein Antibody generated in goat detects immunoglobulin A (alpha chain) from human. Immunoglobulin A (IgA) is an antibody that plays a critical role in mucosal immunity. IgA has two subclasses (IgA1 and IgA2) and can exist in a dimeric form called secretory IgA (sIgA). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays. This product has been overfilled to ensure total recovery of stated quantity.

Safety Information

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL **Diagnostica Vertrieb GmbH** Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

Management System ISO 9001:2015 TÜVRheinland ZERTIFIZIERT

ID 9000019771