

Product Datasheet / Instruction for use



SEC-183783 2023-02-23 1 / 2

Goat F(ab')2 anti-Mouse IgG (H+L)-unconj., MinX Bo,Ho,Hu,Rb,Rt,Sh

General Information

Catalog Number: SEC-183783 Physical State: Liquid (sterile filtered)

Quantity: 0,5 mg Reconstitution Volume:

Reconstitution Buffer:

Shipping Conditions: Wet Ice

Product Expiration Expiration date is one (1) year from date of

opening.

Antibody Host / Format

Host Species: Goat Clonality / Isotype / Clone: polyclonal Ig

Antibody Format: IgG F(ab')2 Fragment Concentration: 1.0 mg/mL by UV absorbance at 280 nm

Conjugation: unconjugated

Maximum Excitation / Emission: /

Specificity

Target Species: Mouse IgG whole molecule

Antibody Specificity: IgG (H+L) This product was prepared from monospecific antiserum by

immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), or Bovine, Horse, Human, Rabbit, Rat, and Sheep Serum Proteins.

Minimal Cross Reactivity: Bovine, Horse, Human, Rabbit, Rat, Sheep

Formulation, Transport and Storage

Storage Buffer: 0.01 M Sodium Phosphate,

0.15 M Sodium Chloride, pH 7.2

Stabilizer: None

Preservatives: 0.01% (w/v) Sodium Azide

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Storage Temperature: 4-8°C

Application Recommendation

Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections),

Immunohistochemistry (frozen sections), Western Blot

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOLDiagnostica Vertrieb GmbH
Leipziger Straße 4
85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com

Your Secondary Antibody Portal Powered by BIOZOL



Management System ISO 9001:2015



www.tuv.com ID 9000019771



Product Datasheet / Instruction for use

SEC-183783 2023-02-23 2 / 2

Dilution: ELISA 1:20.000 - 1:100.000, Immunohistochemistry (IHC) 1:1,000 - 1:5,000, Western Blot (WB) 1:2.000 -

Application Note:

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.

Background Information

F(ab')2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Safety Information

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.







