

Product Datasheet / Instruction for use



# Goat F(ab')2 anti-Mouse IgG (H+L)-Biotin, MinX none

# **General Information**

Catalog Number:	SEC-183799	Physical State:	Lyophilized
Quantity:	0,5 mg	Reconstitution Volume:	500 μL
		<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)
		Shipping Conditions:	at room temperature
		Product Expiration	Expiration date is one (1) year from date of opening.
Antibody Host / I	Format		
Host Species:	Goat	Clonality / Isotype / Clon	e: polyclonal Ig
Antibody Format:	lgG F(ab')2 Fragment	Concentration:	0.5 mg/mL by UV absorbance at 280 nr
		Conjugation:	Biotin
		Maximum Excitation / Emission: /	
Specificity			
Target Species:	Mouse	Immunogen:	Mouse IgG whole molecule
Antibody Specificity: IgG (H+L)			
Antibody Specificit	: <b>y:</b> IgG (H+L)	immunoaffinity chromate beads followed by solid p reactivities, pepsin digest immunoelectrophoresis r	esulted in a single precipitin arc against anti- Mouse IgG and Mouse Serum. No reaction was
Antibody Specificit	: <b>у:</b> IgG (H+L)	immunoaffinity chromato beads followed by solid p reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, I	ography using Mouse IgG coupled to agarose whase adsorption(s) to remove any unwanted cion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- Mouse IgG and Mouse Serum. No reaction was psin or anti-Goat IgG F(c).
	ry: IgG (H+L) nsport and Storage	immunoaffinity chromate beads followed by solid p reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, I observed against anti-Pep	ography using Mouse IgG coupled to agarose whase adsorption(s) to remove any unwanted cion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- Mouse IgG and Mouse Serum. No reaction was psin or anti-Goat IgG F(c).
Formulation, Tra Storage Buffer: 0.	nsport and Storage	immunoaffinity chromato beads followed by solid p reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, I observed against anti-Pep <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -2	ography using Mouse IgG coupled to agarose whase adsorption(s) to remove any unwanted cion and chromatographic separation. Assay b resulted in a single precipitin arc against anti- Mouse IgG and Mouse Serum. No reaction was psin or anti-Goat IgG F(c). : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and
Formulation, Tra Storage Buffer: 0. Phosphate, 0.15 M Stabilizer: 10	nsport and Storage .02 M Potassium	immunoaffinity chromato beads followed by solid p reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, I observed against anti-Pep <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -2 thawing. Centrifuge proc room temperature. This	bgraphy using Mouse IgG coupled to agarose whase adsorption(s) to remove any unwanted ition and chromatographic separation. Assay b resulted in a single precipitin arc against anti- Mouse IgG and Mouse Serum. No reaction was psin or anti-Goat IgG F(c). : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and duct if not completely clear after standing at
Formulation, Train Storage Buffer: 0. Phosphate, 0.15 M Stabilizer: 10 Albumin (BSA) - Imu free	<b>nsport and Storage</b> .02 M Potassium Sodium Chloride, pH 7.2 0 mg/mL Bovine Serum	immunoaffinity chromato beads followed by solid p reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, I observed against anti-Pep <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -2 thawing. Centrifuge proc room temperature. This	bgraphy using Mouse IgG coupled to agarose whase adsorption(s) to remove any unwanted cion and chromatographic separation. Assay b resulted in a single precipitin arc against anti- Mouse IgG and Mouse Serum. No reaction was psin or anti-Goat IgG F(c). : none restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and duct if not completely clear after standing at product is stable for several weeks at 4° C as an

Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use. Management Manufacturer System BIOZOL ISO 9001:2015 Phone +49 (89) 3799 666 6 www.dianova.com Diagnostica Vertrieb GmbH Fax +49 (89) 3799 666 99 TÜVRheinland Your Secondary Antibody Portal Leipziger Straße 4 E-mail info@biozol.de ZERTIFIZIERT www.tuv.com ID 9000019771 85386 Eching Powered by BIOZOL www.biozol.de



#### Product Datasheet / Instruction for use



Dilution: ELISA 1:20.000 - 1:100.000, Immunohistochemistry (IHC) 1:1,000 - 1:5,000, Western Blot (WB) 1:2.000 -1:10.000

#### **Application Note:**

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. This product has been assayed against 1.0 ug of Mouse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:150,000 of the reconstitution concentration is suggested for this product.

### **Background Information**

F(ab')2 Anti-Mouse IgG (H&L) Biotin Antibody generated in goat was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

# **Safety Information**

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

#### Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL **Diagnostica Vertrieb GmbH** Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL



Management System ISO 9001:2015