

Product Datasheet / Instruction for use



Rabbit F(ab')2 anti-Mouse IgG (H+L)-HRPO, MinX Hu

General Information

Catalog Number: SEC-18381	D I	Physical State:	Lyophilized		
Quantity: 0,5 m	g	Reconstitution Volume:	500 μL		
		Reconstitution Buffer:	Restore with deionized water (or equivalent)		
		Shipping Conditions:	at room temperature		
		Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / Format					
Host Species: Rabbit		Clonality / Isotype / Clone: polyclonal Ig			
Antibody Format: IgG F(ab')2	Fragment	Concentration:	0.5 mg/mL by UV absorbance at 280 nr		
		Conjugation:	Horseradish Peroxidase (HRP)		
		Maximum Excitation / Emission: /			
Specificity					
Target Species: Mous	e	Immunogen:	Mouse IgG whole molecule		
Antibody Specificity: IgG (H	I+L)	F(ab')2 Anti-Mouse IgG (H&L) was prepared from monospecific antiseru by immunoaffinity chromatography using Mouse IgG coupled to agaros beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay b immunoelectrophoresis resulted in a single precipitin arc against anti- Peroxidase, anti-Rabbit Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Rabbit IgG F(c) or Huma Serum Proteins.			
		Minimal Cross Reactivity: Human			
Formulation, Transport and	Storage				
Storage Buffer:0.02 M PotassiPhosphate,0.15 M Sodium ChloStabilizer:10 mg/mL BovAlbumin (BSA) - Immunoglobulirfree	ride, pH 7.2 ine Serum	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.			
Preservatives: 0.01% (w/v) G	entamicin	Storage Temperature: 4-8°C			
Sulfate					

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.						
Manufacturer				Management System		
BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015		
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771		



Product Datasheet / Instruction for use



Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

Dilution: ELISA 1:10.000 - 1:50.000, Immunohistochemistry (IHC) 1:500 - 1:2,500, Western Blot (WB) 1:1.000 - 1:10.000

Application Note:

F(ab')2 Anti-Mouse IgG (H&L) has been assayed against 1.0 ug of Mouse IgG in a standard capture ELISA using ABTS (2,2'azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:20,000 of the reconstitution concentration is suggested for this product. Specific conditions for reactivity should be optimized by the end user.

Background Information

F(ab')2 Anti-Mouse IgG (H&L) Peroxidase Antibody generated in rabbit was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays. F(ab')2 Anti-Mouse IgG antibody is ideal for investigators involved in serum component protein research.

Safety Information

This reagent contains Gentamicin Sulfate (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.DO NOT ADD SODIUM AZIDE! Use of Sodium Azide as a preservative will substantially inhibit the enzyme activity of horseradish peroxidase!

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT

Management System ISO 9001:2015