

Product Datasheet / Instruction for use



Rabbit F(ab')2 anti-Mouse IgG (H+L)-RPE, MinX Hu

General Information

Catalog Number: SEC-183813	Physical State:	Lyophilized		
Quantity: 0,5 mg	Reconstitution Volume:	1.0 mL		
	Reconstitution Buffer:	Restore with deionized water (or equivalen		
	Shipping Conditions:	at room temperature		
	Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / Format	·			
Host Species: Rabbit	Clonality / Isotype / Clon	ne: polyclonal Ig		
Antibody Format: IgG F(ab')2 Fragment	Concentration:	0.5 mg/mL by absorbance = 82.0 at 5 nm		
	Conjugation:	RPE (R-phycoerythrin)		
	Maximum Excitation / Emission: 490 nm;545 nm;566 nm / 580 nm			
Specificity	I			
Target Species:MouseAntibody Specificity:IgG (H+L)	Immunogen:	Anti-Mouse IgG was produced by repeated immunization with Mouse IgG whole molecule in rabbit.		
	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Coupling to R-PE was followed by size exclusion chromatography to purify conjugate from unreacted R-PE and antibody. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti- Phycoerythrin, anti-Rabbit Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Rabbit IgG F(c) or Humar Serum Proteins.			
	Minimal Cross Reactivity: Human			
Formulation, Transport and Storage	I			
Storage Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2	Stabilizer:10 mg/mL Bovine Serum Albumin (BSA) -Immunoglobulin and Protease free			
	Preservatives: 0.01% (w/v) Sodium Azide		

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.					
Manufacturer				Management System	
BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99	www.dianova.com Your Secondary Antibody Portal	TÜVRheinland	ISO 9001:2015	
85386 Eching	E-mail info@biozol.de www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771	



Product Datasheet / Instruction for use



after standing at room temperature. Do not freeze after reconstitution.

Store reagent in the dark. Use subdued lighting during handling and

4-8°C

Store vial at 4° C prior to restoration. Restore with deionized water (or equivalent). This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Centrifuge product if not completely clear

Application Recommendation

Application: Flow Cytometry, Immunofluorescence

Dilution: Flow cytometry 1:100 - 1:250, Immunofluorescence 1:100 - 1:250

Application Note:

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 µg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.

Storage Temperature:

incubation of cells prior to analysis.

Background Information

F(ab')2 Mouse IgG (H&L) Antibody Phycoerythrin Conjugated Pre-Adsorbed was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Mouse IgG (H&L) Antibody Phycoerythrin Conjugated Pre-Adsorbed is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Safety Information

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer BIOZOL Diagnostica Vertrieb GmbH

Leipziger Straße 4

85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT

Management System ISO 9001:2015