




# Donkey F(ab')2 anti-Mouse IgG (H+L)-FITC, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Rb,Rt,Sh

# **General Information**

Catalog Number:	SEC-183815	Physical State:	Lyophilized
Quantity:	0,5 ml	Reconstitution Volume:	500 μL
		Reconstitution Buffer:	Restore with deionized water (or equivalent)
		Shipping Conditions:	at room temperature
		Product Expiration	Expiration date is one (1) year from date of opening.
Antibody Host / F	Format		
Host Species:	Donkey	Clonality / Isotype / Clon	e: polyclonal Ig
Antibody Format:	lgG F(ab')2 Fragment	Concentration:	1.0 mg/mL by UV absorbance at 280 nm
		Conjugation:	FITC (Fluoresceinisothiocyanat)
		Maximum Excitation / Emission: 492 nm / 520 nm	
Specificity		I	
Target Species:	Mouse	Immunogen:	Mouse IgG whole molecule
		This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti- Fluorescein, anti-Donkey Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c) or Bovine Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Rabbit, Rat and Shee Serum Proteins.	
Antibody Specificit	<b>y:</b> IgG (H+L)	immunoaffinity chromato beads followed by solid p reactivities, pepsin digest immunoelectrophoresis r Fluorescein, anti-Donkey reaction was observed ag Chicken, Goat, Guinea Pig	ography using Mouse IgG coupled to agarose whase adsorption(s) to remove any unwanted cion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- Serum, Mouse IgG and Mouse Serum. No gainst anti-Pepsin, anti-Donkey IgG F(c) or Bovin
Antibody Specificit	<b>y:</b> IgG (H+L)	immunoaffinity chromato beads followed by solid p reactivities, pepsin digest immunoelectrophoresis r Fluorescein, anti-Donkey reaction was observed ag Chicken, Goat, Guinea Pig Serum Proteins. Minimal Cross Reactivity	bgraphy using Mouse IgG coupled to agarose whase adsorption(s) to remove any unwanted cion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- Serum, Mouse IgG and Mouse Serum. No gainst anti-Pepsin, anti-Donkey IgG F(c) or Bovin
	y: IgG (H+L) nsport and Storage	immunoaffinity chromato beads followed by solid p reactivities, pepsin digest immunoelectrophoresis r Fluorescein, anti-Donkey reaction was observed ag Chicken, Goat, Guinea Pig Serum Proteins. Minimal Cross Reactivity	ography using Mouse IgG coupled to agarose whase adsorption(s) to remove any unwanted cion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- Serum, Mouse IgG and Mouse Serum. No gainst anti-Pepsin, anti-Donkey IgG F(c) or Bovin g, Hamster, Horse, Human, Rabbit, Rat and Shee : Bovine, Chicken, Goat, Guinea Pig, Hamster
Storage Buffer: 0. 0.25 M Sodium Chlo Stabilizer: 10	<b>nsport and Storage</b> 01 M Sodium Phosphate,	immunoaffinity chromato beads followed by solid p reactivities, pepsin digest immunoelectrophoresis r Fluorescein, anti-Donkey reaction was observed ag Chicken, Goat, Guinea Pig Serum Proteins. <b>Minimal Cross Reactivity</b> generally, Horse, Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge proc room temperature. This	ography using Mouse IgG coupled to agarose whase adsorption(s) to remove any unwanted cion and chromatographic separation. Assay by resulted in a single precipitin arc against anti- Serum, Mouse IgG and Mouse Serum. No gainst anti-Pepsin, anti-Donkey IgG F(c) or Bovin g, Hamster, Horse, Human, Rabbit, Rat and Shee : Bovine, Chicken, Goat, Guinea Pig, Hamster





#### Product Datasheet / Instruction for use



### **Application Recommendation**

Application: FLISA, Flow Cytometry, Immunofluorescence

**Dilution:** FLISA 1:10.000 - 1:50.000, Flow cytometry 1:500 - 1:2,500, Immunofluorescence 1:1,000 - 1:5,000

#### **Application Note:**

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

# **Background Information**

F(ab')2 Anti-Mouse IgG (H&L) Fluorescein Antibody generated in donkey was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

# **Safety Information**

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

# Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL **Diagnostica Vertrieb GmbH** Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL



Management System ISO 9001:2015