



# Goat F(ab')2 anti-Rabbit IgG (F(ab')2)-Biotin, MinX Bo,Ho,Hu,Ms,Rt,Sh

## **General Information**

Catalog Number: SE	C-183840	Physical State:	Lyophilized		
Quantity:	0,5 mg	Reconstitution Volume:	500 μL		
		<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)		
		Shipping Conditions:	at room temperature		
		Product Expiration	Expiration date is one (1) year from date of opening.		
Antibody Host / Forn	nat				
Host Species: Go	pat	Clonality / Isotype / Clon	ne: polyclonal Ig		
Antibody Format: IgC	G F(ab')2 Fragment	Concentration:	1.0 mg/mL by UV absorbance at 280 nr		
		Conjugation:	Biotin		
		Maximum Excitation / Emission: /			
Specificity					
Target Species:	Rabbit	Immunogen:	Rabbit IgG F(ab')2 fragment		
Antibody Specificity:	lgG (F(ab')2 Fragment)	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay k immunoelectrophoresis resulted in a single precipitin arc against anti- Biotin, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(ab')2 and Rabbit Serun No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rabbi IgG F(c) or Bovine, Horse, Human, Mouse, Rat and Sheep Serum Proteir			
		reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, F No reaction was observed	ion and chromatographic separation. Assay be esulted in a single precipitin arc against anti- Rabbit IgG, Rabbit IgG F(ab')2 and Rabbit Serum d against anti-Pepsin, anti-Goat IgG F(c), Rabbit		
		reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, F No reaction was observed IgG F(c) or Bovine, Horse,	ion and chromatographic separation. Assay by esulted in a single precipitin arc against anti- Rabbit IgG, Rabbit IgG F(ab')2 and Rabbit Serum d against anti-Pepsin, anti-Goat IgG F(c), Rabbit		
Formulation, Transpo	ort and Storage	reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, F No reaction was observed IgG F(c) or Bovine, Horse,	ion and chromatographic separation. Assay by esulted in a single precipitin arc against anti- Rabbit IgG, Rabbit IgG F(ab')2 and Rabbit Serum d against anti-Pepsin, anti-Goat IgG F(c), Rabbit Human, Mouse, Rat and Sheep Serum Proteins		
Formulation, Transpo Storage Buffer: 0.02 N Phosphate, 0.15 M Sodi	V Potassium	reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, F No reaction was observed IgG F(c) or Bovine, Horse, <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to a contents and freeze at -20	ion and chromatographic separation. Assay be esulted in a single precipitin arc against anti- Rabbit IgG, Rabbit IgG F(ab')2 and Rabbit Serum d against anti-Pepsin, anti-Goat IgG F(c), Rabbit Human, Mouse, Rat and Sheep Serum Proteins : Bovine, Horse, Human, Mouse, Rat, Sheep restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and		
Storage Buffer: 0.02 N Phosphate, 0.15 M Sodi	M Potassium ium Chloride, pH 7.2 g/mL Bovine Serum	reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, F No reaction was observed IgG F(c) or Bovine, Horse, <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge prod room temperature. This	ion and chromatographic separation. Assay b esulted in a single precipitin arc against anti- Rabbit IgG, Rabbit IgG F(ab')2 and Rabbit Serun d against anti-Pepsin, anti-Goat IgG F(c), Rabbit Human, Mouse, Rat and Sheep Serum Protein : Bovine, Horse, Human, Mouse, Rat, Sheep restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and luct if not completely clear after standing at		
Storage Buffer: 0.02 N Phosphate, 0.15 M Sodi Stabilizer: 10 mg Albumin (BSA) - Immun free	M Potassium ium Chloride, pH 7.2 g/mL Bovine Serum	reactivities, pepsin digest immunoelectrophoresis r Biotin, anti-Goat Serum, F No reaction was observed IgG F(c) or Bovine, Horse, <b>Minimal Cross Reactivity</b> Store vial at 4° C prior to contents and freeze at -20 thawing. Centrifuge prod room temperature. This	ion and chromatographic separation. Assay be esulted in a single precipitin arc against anti- Rabbit IgG, Rabbit IgG F(ab')2 and Rabbit Serum d against anti-Pepsin, anti-Goat IgG F(c), Rabbit Human, Mouse, Rat and Sheep Serum Proteins : Bovine, Horse, Human, Mouse, Rat, Sheep restoration. For extended storage aliquot 0° C or below. Avoid cycles of freezing and luct if not completely clear after standing at product is stable for several weeks at 4° C as ar		

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.							
Manufacturer				Management System			
<b>BIOZOL</b> Diagnostica Vertrieb GmbH Leipziger Straße 4	Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de	www.dianova.com	TÜVRheinland	ISO 9001:2015			
85386 Eching	www.biozol.de	Powered by BIOZOL	ZERTIFIZIERT	www.tuv.com ID 9000019771			



#### Product Datasheet / Instruction for use



**Application:** ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections), Immunohistochemistry (frozen sections), Western Blot

Dilution: ELISA 1:20.000 - 1:100.000, Immunohistochemistry (IHC) 1:1,000 - 1:5,000, Western Blot (WB) 1:2.000 - 1:10.000

#### **Application Note:**

F(ab')2 Anti-Rabbit IgG F(ab')2 Biotin Antibody has been assayed against 1.0 ug of Rabbit IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.

#### **Background Information**

F(ab')2 Anti-Rabbit IgG F(ab')2 Biotin Antibody generated in goat detects Rabbit F(ab')2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

### **Safety Information**

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

### Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

BIOZOL Diagnostica Vertrieb GmbH Leipziger Straße 4 85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com Your Secondary Antibody Portal

Powered by BIOZOL

TÜVRheinland ZERTIFIZIERT

Management System ISO 9001:2015