

## Goat F(ab')<sub>2</sub> anti-Rat IgG (Fc)-RPE, MinX Bo,Ho,Hu

### General Information

|                                   |  |
|-----------------------------------|--|
| <b>Catalog Number:</b> SEC-183871 | <b>Physical State:</b> Lyophilized   |
| <b>Quantity:</b> 0,5 mg           | <b>Reconstitution Volume:</b> 1.0 mL   |
|                                   | <b>Reconstitution Buffer:</b> Restore with deionized water (or equivalent)       |
|                                   | <b>Shipping Conditions:</b> at room temperature                                  |
|                                   | <b>Product Expiration:</b> Expiration date is one (1) year from date of opening. |

### Antibody Host / Format

|  |   |
|--|---|
| <b>Host Species:</b> Goat                                | <b>Clonality / Isotype / Clone:</b> polyclonal Ig                     |
| <b>Antibody Format:</b> IgG F(ab') <sub>2</sub> Fragment | <b>Concentration:</b> 0.5 mg/mL by absorbance = 82.0 at 565 nm        |
|  | <b>Conjugation:</b> RPE (R-phycoerythrin)                             |
|  | <b>Maximum Excitation / Emission:</b> 490 nm; 545 nm; 566 nm / 580 nm |

### Specificity

|  |  |
|--|--|
| <b>Target Species:</b> Rat                     | <b>Immunogen:</b> Rat IgG F(c) fragment  |
| <b>Antibody Specificity:</b> IgG (Fc Fragment) | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Coupling to R-PE was followed by size exclusion chromatography to purify conjugate from unreacted R-PE and antibody. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Phycoerythrin, anti-Goat Serum, Rat IgG, Rat IgG F(c) and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rat IgG F(ab) or Bovine, Horse and Human Serum Proteins. |
|  | <b>Minimal Cross Reactivity:</b> Bovine, Horse, Human  |

### Formulation, Transport and Storage

|   |   |
|---|---|
| <b>Storage Buffer:</b> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2         | Store vial at 4° C prior to restoration. Restore with deionized water (or equivalent). This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Centrifuge product if not completely clear after standing at room temperature. Do not freeze after reconstitution. Store reagent in the dark. Use subdued lighting during handling and incubation of cells prior to analysis. |
| <b>Stabilizer:</b> 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |   |
| <b>Preservatives:</b> 0.01% (w/v) Sodium Azide  |   |

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

#### Manufacturer

**BIOZOL**  
Diagnostica Vertrieb GmbH  
Leipziger Straße 4  
85386 Eching

Phone +49 (89) 3799 666 6  
Fax +49 (89) 3799 666 99  
E-mail info@biozol.de  
www.biozol.de

[www.dianova.com](http://www.dianova.com)

Your Secondary Antibody Portal  
Powered by BIOZOL



Management System  
ISO 9001:2015

www.tuv.com  
ID 9000019771



**Storage Temperature:** 4-8°C

### Application Recommendation

**Application:** Flow Cytometry, Immunofluorescence

**Dilution:** Flow cytometry 1:100 - 1:250, Immunofluorescence 1:100 - 1:250

### Application Note:

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain  $1 \times 10^6$  cells in flow cytometry is approximately 1.0 µg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.

### Background Information

F(ab')<sub>2</sub> Anti-Rat IgG F(c) Phycoerythrin Antibody was generated in goat and detects specifically Rat IgG F(c). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

### Safety Information

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

### Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

#### Manufacturer

**BIOZOL**  
Diagnostica Vertrieb GmbH  
Leipziger Straße 4  
85386 Eching

Phone +49 (89) 3799 666 6  
Fax +49 (89) 3799 666 99  
E-mail info@biozol.de  
www.biozol.de

[www.dianova.com](http://www.dianova.com)

Your Secondary Antibody Portal  
Powered by BIOZOL



Management System  
ISO 9001:2015

www.tuv.com  
ID 9000019771

