

Product Datasheet / Instruction for use



SEC-183942 2023-02-23 1 / 2

# Goat Fab anti-Hamster generally IgG (H+L)-Biotin, MinX none

### **General Information**

Catalog Number: SEC-183942 Physical State: Lyophilized

Quantity: 1 mg Reconstitution Volume: 1.0 mL

**Reconstitution Buffer:** Restore with deionized water (or equivalent)

**Shipping Conditions:** at room temperature

**Product Expiration** Expiration date is one (1) year from date of

opening.

# **Antibody Host / Format**

Host Species: Goat Clonality / Isotype / Clone: polyclonal Ig

Antibody Format: IgG Fab Fragment Concentration: 1.0 mg/mL by UV absorbance at 280 nm

Conjugation: Biotin

Maximum Excitation / Emission: /

# **Specificity**

Target Species: Hamster generally Immunogen: Golden Syrian Hamster IgG whole molecule

Antibody Specificity: IgG (H+L)

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Golden Syrian Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin and anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c).

Minimal Cross Reactivity: none

# Formulation, Transport and Storage

**Storage Buffer:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Stabilizer:** 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease

free

**Preservatives:** 0.01% (w/v) Sodium Azide

**Application Recommendation** 

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Storage Temperature:** 4-8°C

Application: ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections),

Immunohistochemistry (frozen sections), Western Blot

This product is for In Vitro experimental use only. Not for Therapeutic or Diagnostic use.

Manufacturer

**BIOZOL**Diagnostica Vertrieb GmbH
Leipziger Straße 4
85386 Eching

Phone +49 (89) 3799 666 6 Fax +49 (89) 3799 666 99 E-mail info@biozol.de www.biozol.de

www.dianova.com

Your Secondary Antibody Portal Powered by BIOZOL



Management System ISO 9001:2015



www.tuv.com ID 9000019771



Product Datasheet / Instruction for use



SEC-183942 2023-02-23 2 / 2

Dilution: ELISA 1:200.000, Immunohistochemistry (IHC) 1:1,000 - 1:5,000, Western Blot (WB) 1:2.000 - 1:10.000

#### **Application Note:**

Suitable for immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.

### **Background Information**

Fab Anti-Golden Syrian Hamster IgG (H&L) Antibody generated in goat detects immunoglobulin g from hamster, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. This Fab Anti-Golden Syrian Hamster IgG Antibody is conjugated to Biotin.

## Safety Information

This reagent contains sodium azide (a poisonous and hazardous substance) as preservative. Although this concentration is not regarded as dangerous to health, appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. Standard Laboratory Practices should be followed when handling this material.

#### Disclaimer

This product is warranted to perform in conformance with product specifications and to be free from defects in material and workmanship. Products are supplied for research use only. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated, this warranty is limited to one year from date of shipment when the product is subjected to normal, proper and intended usage. Buyer's exclusive remedy for non-conforming products during the warranty period is limited to replacement of or refund for the non-conforming product(s). There is no obligation to replace products as the result of accident or disaster or inappropriate use of the products in a manner for which they were not designed, or improper storage and handling of the products.







