Beratung und Vertrieb 🗇 🖓 Fon + 49 (0) 40 45 067 0 • Fax + 49 (0) 40 450 67 490 • info@dianova.de • www.dianova.de • Warburgstr. 45 • 20354 Hamburg



Instructions For Use			
LGB-IFU			
Rev. Date: Aug. 12, 2011	Revision: 1	Page 1 of 1	
20.9250 Tol. (425) 755.0949 Eav. (425) 755.0015 unuu soutok som			

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

Light Green S.F. Yellowish Solution

Description:	Light Green S.F. Yellowish Solution is intended for use in histological applications as a general counterstain. When used correctly, various shades of green can be obtained to aid in visualization of tissue components.		
	Cytoplasm: Nuclei:	Light Green Light Green/Blue	
Uses/Limitations:	Not to be taken internally. For In-Vitro Diagnostic use only. Histological applications. Do not use past expiration date. Use caution when handling reagents. Non-Sterile.		
Availability:	<u>ltem #</u> LGB500 LGB999 LGB-20000	<u>Volume</u> 500 ml 1000 ml 20 liters	
Storage:	Store at 18-25 ℃.		
Precautions:	Avoid contact with skin and eyes. Harmful if swallowed. Follow all Federal, State, and local regulations regarding disposal.		

Procedure:

- 1. Deparaffinize sections if necessary and hydrate to distilled water.
- 2. Rinse slide in distilled water.
- 3. Stain slide in Light Green Solution for 2 minutes.
- 4. Rinse 1 time in Distilled/Deionized water.
- 5. Dehydrate quickly through graded alcohols.
- 6. Clear, and mount in synthetic resin.



Doc: IFU-Template18-25rev2

ScyTek Laboratories, Inc. 205 South 600 West Logan, UT 84321 U.S.A.



EC REP EmergoEurope (31)(0) 70 345-8570 Molsnstraat 15 2513 BH Hague, The Netherlands